CUMULATIVE AUTHOR INDEX
Journal of Medical Entomology
Volume 37, 2000

A

Abbassy, M. M., 393
Abel, I. D. S., 714
Abro, G. H., 534
Agnew, P., 732
Agui, N., 945
Alekseev, A. N., 9
Ali, Z. B., 787
Almirón, W. R., 16
Alto, B. W., 364
Alzogaray, R. A., 308, 957
Amerasinghe, P. H., 89
Anderson, J., 454
Anderson, S., 864
Anderson, S. H., 612
Ansari, M. A., 201
Antoniou, M., 346
Apperson, C. S., 40
Arrivillaga, J., 325

B

Back, C., 29
Bahia Labruna, M., 774, 777
Baimai, V., 340
Baisley, K. J., 53, 340
Bangs, M. J., 840
Baraka, O. Z., 547
Barbanti Duarte, J. M., 979
Barriga, O. O., 601
Barros-Battesti, D. M., 820
Barth, J. F., 581
Beavers, G. M., 393
Bechara, G. H., 979
Beebe, N. W., 153, 840
Beier, J. C., 246, 528
Belden, R. P., 58
Bella, A., 281
Belwood, J. J., 445
Benabdennbi, I., 289
Bergman, D. K., 885
Bergoin, M., 802
Berkebile, D. R., 968
Bhaskaran, G., 503
Biggerstaff, B. J., 559
Blackwell, A., 222
Boakye, D. A., 29, 547
Bobian, R. J., 58, 65
Bonder, M. A., 271
Borges, E. C., 872
Boursot, P., 787
Boyce, K., 507
Boyce, W. M., 791
Boyd, A. M., 660
Brakefield, P. M., 29
Bram, R. A., 1
Brannon, E. S., 559
Brazil, R. P., 860
Brewer, M., 16
Brooks, C. J. W., 222
Bryan, J. H., 153
Buczek, A., 807
Burge, R., 919
Burkhart, C. G., 170
Burkhart, C. N., 170
Burkot, T. R., 467

C

Cabaret, J., 335
Cadiergues, M.-C., 634
Calampa, C., 835
Campbell, C. L., 784
Campos, R. E., 385
Cantaloube, B., 634
Carlson, D. A., 308, 957
Carroll, J. F., 472
Castellón, E. G., 162
Cavagnari, B. M., 938
Cedeno, V., 802
Cepeda-Palacios, R., 239
Chae, J.-s., 619
Chaniotis, B., 346
Chapman, H. F., 736
Charlab, R., 743
Chatti, N., 787
Chebret, M., 769
Chen, W.-J., 108
Chiles, R. E., 250, 259, 507
Chiou, L.-Y., 108
Chirebvu, E., 286
Chirico, J., 968
Chow-Shaffer, E., 492
Chuang, W.-L., 108
Chuang, Y.-Y., 205
Cilek, J. E., 439
Clark, G. G., 77, 89
Clerici, P. T. S., 714
Coleman, J., 102
Coleman, R. E., 581, 835
Colwell, D. D., 854
Conn, J. E., 754
Cooper, L. A., 815
Cooper, R. D., 153, 840
Copeland, R., 581
Crabtree, M. B., 554
Crandell, M., 454
Cribb, B. W., 316
Cueto, G. A. M., 721
Cupp, E. W., 416, 743
Cupp, M. S., 416
Cusack, R., 507

D

Da Cunha Ramos, H., 962
Daniels, T. J., 357
Danishefsky, K., 102
Darwish, Z. E. A., 393
Daugherty, M. P., 364
Davila, G., 541
Davis, B. S., 559
Day, J. F., 626, 977
de Aluja, A. S., 489
Dean, S., 503
Debboun, M., 177, 919
De Benedictis, J., 492
De Campos Pereira, M., 979
de Castro Gomes, A., 820
de Melo Ximenes, M. F. F., 162
Derock, E., 619
de Souza, M. F., 162

Deunff, J., 445
Díez, N., 27
Díez, P., 27
Di Luca, M., 281
Diotaiuti, L., 373, 872
Do, K.-A., 146
Dohm, D. J., 435, 835
Dolan, M. C., 766
Dong, C.-F., 108
Drey, C., 761
Dubinina, H. V., 9
Dujardin, J. P., 872
Dunton, R. F., 467
Durden, L. A., 276

E

Edman, J. D., 77, 89, 701
Eldridge, B. F., 250, 259, 507
Elms, S. L., 909
Engelthaler, D. M., 422
Eraso, E., 35
Ewald-Hamm, D., 228

F

Fairley, T. L., 754
Falco, R. C., 357
Fampa, P., 802
Fang, Q. Q., 276
Farrell, B. D., 791
Farrow, R. A., 797
Federici, B. A., 401
Feliciangeli, M. D., 325
Fernandez, R., 835
Ferreira Homem, V. S., 774, 777
Ferreira Neto, J. S., 774, 777
Fielden, L., 979
Fielden, L. J., 761
Figureau, C., 335
Fish, D., 357
Flanary, B. E., 950
Foley, D. H., 933
Foley, P. N., 846
Fournier, P. E., 707
Foy, B. D., 528
Franc, M., 634
Freitas, R. A., 162
Fryauff, D. J., 393
Fujioka, K. K., 295

G

Gage, K. L., 422
Garcia, A., 541
Gemmell, M., 489
George, J. E., 1, 588, 878
Georghiou, G. P., 534
Gerez De Burgos, N. M., 689, 938
Gerry, A. C., 675
Ghosh, K., 575
Gimnig, J. E., 902
Giovannangeli, S., 281
Gleiser, R. M., 21
Glenister, C. S., 929
Goldberg, M., 761
Golenda, C., 919
Gonzalez-Ceron, L., 331
Gooding, R. H., 897
Gordon, J. C., 601
Gordon, S. W., 581
Gorla, D. E., 21
Gregory, L. M., 571
Grillet, M. E., 231
Grimstad, P., 454
Grobbelaar, A. A., 467
Grogan, W. L., 65
Grunewald, J., 748
Guarneri, A. A., 373
Guisantes, J. A., 35
Gupta, R., 182
Gutierrez, A. G., 349
Gutierrez, D., 507
Gutierrez-Palomino, P., 210
Guy, J. S., 480

H

Haddad, N., 289
Haller, J. F., 938
Haramis, L., 454
Haussy, C., 732
Hawley, W. A., 492
Heinemann, M. B., 774, 777
Hernandez, J. E., 331
Hernández-Rodríguez, S., 210
Herrero, M. V., 216
Higazi, T. B., 547
Hill, B. R., 507
Hogsette, J. A., 957
Holbrook, F. R., 65
Holman, M. S., 126
Holt, J. S., 846
Hourcq, P., 634
Hsu, M. H., 828
Hu, R., 198
Hughes, J. M., 736
Hung, R. W., 121
Hutchinson, M. J., 228
Hynes, W. L., 265

I

Ilango, K., 653
Irving, R. P., 595

J

Jacobs, D. E., 228
Jacobson, R. L., 319
Jaenson, T. G. T., 973
Jamnongluk, W., 53
Jerônimo, S. M. B., 162
Jimenez, A. E., 216
Jittapalapong, S., 601
Johns, R., 265
Johnson, E., 619
Jones, C. J., 408
Jones, J. W., 835
Jordan, R. A., 121
Juliano, S. A., 364, 950
Jupp, P. G., 467

K

Kamimura, K., 637
Karabatsos, N., 559
Karhu, R., 864
Karhu, R. R., 612
Karpowicz, J. C., 909
Kaufman, P. E., 929
Kay, B. H., 146, 380, 660, 736, 797, 846
Keep, L., 919
Keirans, J. E., 979
Kelley, S. T., 791
Kemp, A., 467
Killeen, G. F., 528
Kitron, U., 454
Kitron, U. D., 408
Kittayapong, P., 53, 77, 89, 340
Klein, T. A., 835, 919
Klun, J. A., 177, 182
Knerl, K. M., 909

Knipling, E. F., 645
Kobayashi, M., 945
Kollars, T. M., Jr., 640
Kovaltchouk, J. G., 295
Kramer, L. D., 250, 259
Kramer, V. L., 909
Kristensen, P. K., 984
Ksiazek, T. G., 467
Kuramochi, K., 461

L

Labruna, M. B., 714
Lacombe, E. H., 126, 766
Lagnel, J., 787
Lancaster, M., 454
Lane, R. P., 653
Lane, R. S., 484
Lardeux, F., 769
La Torre, J. L., 271
Lawler, S. P., 619
Lawyer, P. G., 435
Lee, J.-H., 187, 194
Leger, N., 289
Leman, P. A., 467
Lemeilleur, C. A., 588
Leopold, R. A., 968
Lerdthusnee, K., 893
Lien, J.-C., 780
Lin, C.-C., 780
Lin, C.-S., 205
Lindblade, K. A., 664
Liu, H. Q., 559
Long, S. J., 929
Lonning, S. M., 784
López, C., 27
Lord, C. C., 977
Lorenz, L. H., 77, 89
Lothrop, H. D., 507
Lounibos, L. P., 385
Lubelczyk, C., 126
Ludueña Almeida, F. F., 16
Lundstrom, J. O., 507
Lyons, S. A., 846
Lysyk, T. J., 299

M

Ma, D., 182
Mackey, A. J., 246
Madigan, J. E., 619
Mahmoud, B. M., 547
Malik, J. I., 534
Mangiafico, J. A., 965
Manokore, V., 286
Manweiler, S. A., 885
Martínez, J., 35
Martinez, M. J., 489
Martinez, V. M., 250, 259, 507
Martínez-Moreno, F. J., 210
Marzouk, A. S., 393
Matushima, E. R., 979
Mboera, L. E. G., 172
McKinnon, C. N., 65
McPherson, K. R., 919
McPherson, M., 114
Mdira, K. Y., 172
Meier, K., 503
Mejlon, H. A., 973
Meola, R., 503
Meola, R. W., 141
Mialhe, E., 802
Michalakis, Y., 732
Miller, B. R., 554
Miller, J. A., 588, 878
Miller, L. H., 802
Miller, M. J., 909
Miyagi, I., 554
Mohamed, F. S. A., 393
Moll, R. M., 893
Moncayo, A. C., 701, 893
Moore, C. B., 427
Moore, C. G., 559
Mora, J., 965
Moribayashi, A., 726
Morohashi, M., 637
Morrison, A. C., 77, 89
Morrondo, P., 27
Mukhopadhyay, J., 575
Mukhtar, M. M., 547
Mullens, B. A., 675
Murahwa, F. C., 286
Muscio, O., 271

N

Nalyanya, G., 427
Nasci, R. S., 559
Navajas, M., 787
Navarrete, I., 210
Nazario Bonoldi, V. L., 820
Nelson, M., 541
Nieves, E., 134
Nunamaker, R. A., 784

O

O'Brien, J. J., 349
O'Connor, M., 854
O'Guinn, M., 435
Olesen, J. B., 595
Oliver, J. H., Jr., 276
Olson, M. A., 439
O'Neill, S. L., 53, 340
Oviedo, M., 325

P

Palacios, R., 35
Panadero, R., 27
Panella, N. A., 559
Parker, B. M., 695
Paz, A., 27
Pearson, R. D., 162
Pereira, M. H., 373
Peréz De León, A. A., 784
Pérez-Eid, C., 335
Perich, M. J., 541
Perrotey, S., 289
Pesce, M., 281
Pesson, B., 289
Pichon, B., 335
Pickett, J. A., 172
Picollo, M. I., 721
Piesman, J., 766
Pimenta, P. F. P., 134
Pinger, R. R., 595
Platt, K. B., 187, 194, 965
Pound, J. M., 588, 878
Prescott, R. J., 222
Pusterla, N., 619

R

Raineri, W., 281
Ramachandra, N., 102
Ramey, R. R., 791
Rand, P. W., 126
Randolph, M. P., 349
Rangel, Y., 325
Raoult, D., 707
Rasa, C. G., 141
Razdan, R. K., 201
Rechav, Y., 761, 979
Řeháček, J., 707
Reina, D., 210
Reisen, W. K., 250, 259, 507

Renaud, T. M., 754
Rey, A., 281
Rhein, S. L. V., 950
Ribeiro, H., 962
Ribeiro, J. M. C., 743
Ritchie, S. A., 153, 736
Roake, W., 528
Rocha, O. J., 965
Rodriguez, M. H., 331
Rojas, J. C., 216
Romanha, A. J., 872
Romi, R., 281
Romoser, W. S., 893
Rowley, W. A., 187, 194, 198, 965
Rowton, E. D., 435, 743
Russell, B. M., 846
Rutz, D. A., 929
Ryan, P. A., 146, 846

S

Said, K., 787
Samuel, W. M., 114
Santillan, F. V., 331
Santomé, J. A., 938
Saraiva, E., 802
Sardelis, M. R., 835
Sasaki, T., 945
Sattabongkot, J., 581
Saul, A., 380, 933
Saviskas, B., 295
Sawabe, K., 726
Scaraffia, P. Y., 689, 938
Schal, C., 427, 924
Schelotto, G., 21
Schenker, R., 141
Schlein, Y., 319
Schmidtmann, E. T., 58, 65
Schoeler, G. B., 885
Schofield, C. J., 872
Scholl, P. J., 210, 239
Schulze, T. L., 121
Schumaker, T. T. S., 714
Schwartz, I., 102
Scodeller, E. A., 271
Scott, T. W., 77, 89, 492, 507, 815
Scoulika, E., 346
Sekeyová, Z., 707
Seki, T., 637
Shahabuddin, M., 528
Shaio, M.-F., 641
Shirai, Y., 637
Shoop, W. L., 571
Shostak, A. W., 114
Shoukry, M. A., 393
Silva, O. S., 748
Sina, B., 492
Sina, B. J., 815
Sithiprasasna, R., 640
Smith, R. P., Jr., 126
Soares, M. J., 860
Solberg, V. B., 919
Somboon, P., 476
Sommer, C., 984
Sonenshine, D. E., 265
Spiegel, C. N., 860
Spyridaki, I., 346
Stark, L. M., 626
Steelman, C. D., 645
Steiner, F. E., 595
Stich, R. W., 601
Strickman, D., 77, 89, 919
Stringham, S. M., 480
Stromdahl, E. Y., 349
Stuart, A. E., 222
Swanepoel, R., 467
Swanson, J., 454
Szabó, M. P. J., 979

T

Tabachnick, W. J., 65
Takagi, M., 476
Takken, W., 172
Tälleklint-Eisen, L., 484
Teng, H.-J., 40
Terenius, O., 973
Tetreault, G. E., 393
Theis, J. H., 295
Thompson, M. A., 909
Toma, T., 554
Torres, E. P., 933
Touré, A. M., 246
Trueba, G., 965
Tsuda, Y., 476
Tuno, N., 476
Turell, M. J., 435, 581, 701, 815, 835
Turner, A., 541

U

Ung, A., 769
Unnasch, T. R., 547

V

van den Hurk, A. F., 153, 736
Vann, C. N., 595
Vargas, F., 216
Vassallo, M., 335
Vassena, C. V., 721
Vaughan, T. J., 528
Vernetti, M., 721

W

Walker, E. D., 664
Walton, W. E., 401
Wang, C.-H., 205
Wang, Z. X., 246
Waterson, D. G. E., 840
Watson, D. W., 480, 924
Watson, T. M., 380
Watts, D. M., 835
Weng, M.-H., 641, 780
Whitaker, J. O., Jr., 445
Whitlock, J. E., 276
Wikel, S. K., 885
Wilkerson, R. C., 919
Williams, A., 528
Williams, G. M., 153
Wilson, M. D., 547
Wilson, M. E., 162
Wilson, M. L., 664
Wirth, M. C., 401, 534
Wirtz, R. A., 331, 581, 919
Wittum, T. E., 601
Wright, S. A., 909
Wright, S. E., 507
Wu, W. J., 828

Y

Yao, C.-W., 641, 780
Yeh, C.-C., 205
Yoshinari, N. H., 820
Young, J. F., 909

Z

Zakson-Aiken, M., 571
Zaldivar, M. E., 965
Zanasi, F., 281
Zerba, E. N., 721
Zhang, D., 416
Zhou, H., 77
Zurek, L., 924

CUMULATIVE SUBJECT INDEX
Journal of Medical Entomology
Volume 37, 2000

A

abundance,
  accuracy, 299
  bias, 299
  Haematobia irritans, 299
  precision, 299
  sampling, 299
abundance variation,
  Aedes albifasciatus, 16
  autumn–winter, 16
  gonotrophic status, 16
acaracide,
  Amblyomma variegatum, 645
  Ixodes scapularis, 645
  Ixodiphagus hookeri, 645
  parasitoid augmentation, 645
acaricide, 588, 878
  Amblyomma americanum, 588, 878
  amitraz, 588, 878
  lone star tick, 588, 878
  self-treatment, 588, 878
acaricides,
  Amblyomma americanum, 121
  control, 121
  Ixodes scapularis, 121
  Lyme disease, 121
accuracy,
  abundance, 299
  bias, 299
  Haematobia irritans, 299
  precision, 299
  sampling, 299
acquired resistance,
  concanavalin A, 885
  Ixodes scapularis, 885
  lipopolysaccharide, 885
  lymphocytes, 885
  salivary gland, 885
adaptations,
  behavior, 114
  Dermacentor albipictus, 114
  host-finding, 114
  larvae, 114
  transmission, 114
adult,
  cryopreservation, 968
  primary screwworm, 968
  pupa, 968
  survival, 968
aedeagal filaments,
  arms race, 653
  coadaptation, 653
  Phlebotomine sand flies, 653
  spermathecal ducts, 653
Aedes abserratus,
  Aedes hexodontus, 902
  Aedes punctor, 902
  allozymes, 902
  genetic clines, 902
  larval morphology, 902
Aedes aegypti. 77, 89, 364, 492, 541, 641, 660, 846, 965
  Aedes albopictus, 364
  Aedes notoscriptus, 660
  antibodies, 965
  antibody effect, 641
  anti-dengue serum, 641
  arbovirus, 846
  Australia, 846
  Barmah Forest virus, 660
  bats, 965
  blood feeding, 492
  Culex annulirostris, 660
  Culex quinquefasciatus, 660
  Culex sitiens, 660
  culicidae, 541
  dengue virus, 641, 965
  DNA fingerprinting, 492
  histology, 89
  insect container communities, 364
  invertebrate carcasses, 364
  Mesocyclops, 846
  mosquito, 846
  multiple blood feeding, 89
  neutralizing, 965
  population dynamics, 77
  Puerto Rico, 77, 89
  replacement series, 364
  resting behavior, 541
  subterranean, 846
  Thailand, 77, 89
  ultra-low volume insecticides, 541

virus isolation methods, 641
Aedes albifasciatus, 16, 21
abundance variation, 16
Argentina, 21
autumn–winter, 16
gonotrophic status, 16
population dynamics, 21
Aedes albopictus, 40, 281, 364, 454, 637, 701
Aedes aegypti, 364
Aedes triseriatus, 40, 454
alightment, 637
Anopheles quadrimaculatus, 701
bite, 637
competition, 454
Coquillettidia perturbans, 701
degree-day, 40
eastern equine encephalomyelitis virus, 701
immature development, 40
insect container communities, 364
invasion, 454
invertebrate carcasses, 364
La Crosse virus, 454
larval development, 281
metallic copper, 281
mosquito attractant, 637
mosquito control, 281
mosquito repellent, 637
oviposition, 454
proboscis amputation, 637
replacement series, 364
survivorship, 701
Aedes canadensis,
Aedes triseriatus, 559
habitat structure, 559
La Crosse encephalitis virus, 559
mosquito ecology, 559
Aedes communis,
AI3-37220, 919
deet, 919
piperidine, 919
repellents, 919
Simulium venustum, 919
Aedes hexodontus,
Aedes abserratus, 902
Aedes punctor, 902
allozymes, 902
genetic clines, 902
larval morphology, 902
Aedes notoscriptus, 380, 660
Aedes aegypti, 660
Australia, 380
Barmah Forest virus, 380, 660
Culex annulirostris, 660
Culex quinquefasciatus, 660
Culex sitiens, 660
dispersal, 380
mark-release-recapture, 380
Ross River virus, 380
Aedes polynesiensis,
age-grading, 769
Culex quinquefasciatus, 769
fluorescence, 769
pteridines, 769
spectrofluorometry, 769
Aedes punctor,
Aedes abserratus, 902
Aedes hexodontus, 902
allozymes, 902
genetic clines, 902
larval morphology, 902
Aedes sollicitans,
Aedes taeniorhynchus, 695
Dirofilaria immiti, 695
dog heartworm, 695
mosquitoes, 695
vector competence, 695
Aedes taeniorhynchus,
Aedes sollicitans, 695
Dirofilaria immiti, 695
dog heartworm, 695
mosquitoes, 695
vector competence, 695
Aedes triseriatus, 40, 454, 559, 950
Aedes albopictus, 40, 454
Aedes canadensis, 559
Ascogregarina barretti, 950
competition, 454
degree-day, 40
habitat drying, 950
habitat structure, 559
immature development, 40
invasion, 454
La Crosse encephalitis virus, 559
La Crosse virus, 454
mosquito ecology, 559
oviposition, 454
prevalence intensity of infection, 950
age-grading,
Aedes polynesiensis, 769

Culex quinquefasciatus, 769
fluorescence, 769
pteridines, 769
spectrofluorometry, 769
AI3-37220,
Aedes communis, 919
deet, 919
piperidine, 919
repellents, 919
Simulium venustum, 919
airborne action,
Anopheles culicifacies, 201
Culex quinquefasciatus, 201
excito-repellency action, 201
insecticide-treated mosquito nets, 201
killing action, 201
alightment,
Aedes albopictus, 637
bite, 637
mosquito attractant, 637
mosquito repellent, 637
proboscis amputation, 637
allozymes,
Aedes abserratus, 902
Aedes hexodontus, 902
Aedes punctor, 902
genetic clines, 902
larval morphology, 902
altitude,
Anopheles, 664
epidemic, 664
forecasting methods, 664
malaria, 664
sample size, 664
Amblyomma americanum, 121, 276, 349, 439, 588, 595, 878
acaricide, 588, 878
acaricides, 121
Amblyomma maculatum, 439
amitraz, 588, 878
barrier islands, 276
coastal Georgia, 276
control, 121
Dermacentor variabilis, 439
Ehrlichia chaffeensis, 276, 349, 595
human monocytic ehrlichiosis, 276, 349, 595
indirect immunofluorescence assay, 595
Ixodes scapularis, 121, 439
lone star tick, 588, 878
Lyme disease, 121
Maryland, 349
polymerase chain reaction, 349, 595
prevalence, 276
self-treatment, 588, 878
tick ecology, 439
Amblyomma auricularium,
armadillo, 977
Dasypus novemcinctus, 977
Florida, 977
Amblyomma cajennense,
Amblyomma oblongoguttatum, 774
Amblyomma ovale, 774
Brazil, 774
dogs, 774
ticks, 774
Amblyomma maculatum, 439, 761
Amblyomma americanum, 439
capillary feeding, 761
Dermacentor variabilis, 439
Ixodes scapularis, 439
Ixodid ticks, 761
production of aggregation-attachment pheromones, 761
tick ecology, 439
Amblyomma oblongoguttatum, 774, 777
Amblyomma cajennense, 774
Amblyomma ovale, 774
Brazil, 774
dogs, 774
gynandromorphism, 777
ticks, 774, 777
Amblyomma ovale,
Amblyomma cajennense, 774
Amblyomma oblongoguttatum, 774
Brazil, 774
dogs, 774
ticks, 774
Amblyomma variegatum,
acaracide, 645
Ixodes scapularis, 645
Ixodiphagus hookeri, 645
parasitoid augmentation, 645
amitraz, 588, 878
acaricide, 588, 878
Amblyomma americanum, 588, 878
lone star tick, 588, 878
self-treatment, 588, 878
Androctonus,
genetic diversity, 787

ITS, 787
ribosomal DNA, 787
scorpions, 787
Angola,
Anopheles squamosus, 962
Coelomomyces numularius, 962
new species, 962
anomalies,
embryonic development, 807
Hyalomma marginatum, 807
teratology, 807
Anopheles,
altitude, 664
epidemic, 664
forecasting methods, 664
malaria, 664
sample size, 664
Anopheles albimanus,
phenotypes, 331
Plasmodium vivax, 331
susceptibility, 331
Anopheles aquasalis,
Anopheles oswaldoi, 231
environmental factors, 231
larval habitats, 231
malaria, 231
spatial temporal distribution, 231
Anopheles arabiensis,
Anopheles gambiae complex, 286
DDT, 286
susceptibility, 286
synthetic pyrethroids, 286
Anopheles clowi,
Anopheles punctulatus group, 840
Papua New Guinea, 840
Anopheles culicifacies,
airborne action, 201
Culex quinquefasciatus, 201
excito-repellency action, 201
insecticide-treated mosquito nets, 201
killing action, 201
Anopheles dispar,
Anopheles greeni, 933
Anopheles maculatus, 933
ITS2, 933
Philippines, 933
polymerase chain reaction-restriction fragment-length polymorphism, 933
Anopheles farauti,
Australia, 153
malaria, 153
seasonal abundance, 153
sibling species, 153
Anopheles flavirostris,
Anopheles minimus, 476
cibarial armature, 476
hybridization, 476
Indonesia, 476
Anopheles gambiae, 246, 528
Anopheles stephensi, 246
antibiotics, 246
bacteria, 246
Monoclonal antibody, 528
phage display, 528
scFv, 528
screening, 528
transmission blocking, 528
Anopheles gambiae complex,
Anopheles arabiensis, 286
DDT, 286
susceptibility, 286
synthetic pyrethroids, 286
Anopheles greeni,
Anopheles dispar, 933
Anopheles maculatus, 933
ITS2, 933
Philippines, 933
polymerase chain reaction-restriction fragment-length polymorphism, 933
Anopheles maculatus,
Anopheles dispar, 933
Anopheles greeni, 933
ITS2, 933
Philippines, 933
polymerase chain reaction-restriction fragment-length polymorphism, 933
Anopheles minimus,
Anopheles flavirostris, 476
cibarial armature, 476
hybridization, 476
Indonesia, 476
Anopheles oswaldoi,
Anopheles aquasalis, 231
environmental factors, 231
larval habitats, 231
malaria, 231
spatial temporal distribution, 231
Anopheles punctipennis,

COI gene, 754
mitochondrial DNA, 754
population structure, 754
Anopheles punctulatus group,
Anopheles clowi, 840
Papua New Guinea, 840
Anopheles quadrimaculatus,
Aedes albopictus, 701
Coquillettidia perturbans, 701
eastern equine encephalomyelitis virus, 701
survivorship, 701
Anopheles squamosus,
Angola, 962
Coelomomyces numularius, 962
new species, 962
Anopheles stephensi, 177, 182, 246, 726
Anopheles gambiae, 246
antibiotics, 246
bacteria, 246
Culex pipiens molestus, 726
Culex pipiens pallens, 726
1-[3-cyclohexen-1-ylcarbonyl]-2-methylpiperidine, 177, 182
1-[3-cyclohexen-1-ylcarbonyl] piperidine, 177, 182
fatty acids, 726
lipids, 726
N,N-diethyl-3-methylbenzamide, 177
new module, 177
repellents, 182
stereoisomers, 182
yolk deposition, 726
antenna,
Indian sand flies, 575
larvae, 575
mouthparts, 575
Phlebotomus, 575
Sergentomyia, 575
antibiotics,
Anopheles gambiae, 246
Anopheles stephensi, 246
bacteria, 246
antibodies,
Aedes aegypti, 965
bats, 965
dengue virus, 965
neutralizing, 965
antibody, 250, 259
Culex tarsalis, 250
house finch, 259
infection, 259
mosquito blood feeding, 250
St. Louis encephalitis virus, 250, 259
viremia, 250, 259
western equine encephalomyelitis virus, 250, 259
antibody effect,
Aedes aegypti, 641
anti-dengue serum, 641
dengue virus, 641
virus isolation methods, 641
anti-dengue serum,
Aedes aegypti, 641
antibody effect, 641
dengue virus, 641
virus isolation methods, 641
antifeedant,
Culicoides impunctatus, 222
6-methylsalicylic acid, 222
repellent, 222
salicylic acid, 222
salicyluric acid, 222
antimicrobial peptides,
Borrelia burgdorferi, 265
Dermacentor variabilis, 265
phagocytosis, 265
antithrombin,
blood-feeding, 416
Hematobia irritans, 416
aquatic ecology,
aquatic insects, 58
biting midges, 58
soil chemistry, 58
vector distribution, 58
aquatic insect host,
Ehrlichia risticii, 619
Potomac horse fever, 619
trematode vector, 619
aquatic insects,
aquatic ecology, 58
biting midges, 58
soil chemistry, 58
vector distribution, 58
arboviral transmission,
mosquito-borne encephalitis, 626
St. Louis encephalitis virus, 626
vector infection rates, 626
arbovirus,
Aedes aegypti, 846
Australia, 846

Mesocyclops, 846
mosquito, 846
subterranean, 846
arbovirus infection,
Culiseta melanura, 815
eastern equine encephalomyelitis virus, 815
mosquito mortality, 815
virulence, 815
Argentina,
Aedes albifasciatus, 21
population dynamics, 21
armadillo,
Amblyomma auricularium, 977
Dasypus novemcinctus, 977
Florida, 977
Armigeres subalbatus, 53, 108
cytoplasmic incompatibility, 53
ecology, 108
Japanese encephalitis virus, 108
rice-free, 108
Taiwan, 108
Wolbachia, 53
arms race,
aedeagal filaments, 653
coadaptation, 653
Phlebotomine sand flies, 653
spermathecal ducts, 653
arrestant,
avoidance, 472
deer, 472
Ixodes scapularis, 472
kairomone, 472
ticks, 472
arthropod pest introductions,
arthropod-borne diseases, 1
nonindigenous arthropods, 1
arthropod-borne diseases,
arthropod pest introductions, 1
nonindigenous arthropods, 1
artificial medium,
bacteria, 924
house fly, 924
intestinal tract, 924
larval development, 924
Ascogregarina barretti,
Aedes triseriatus, 950
habitat drying, 950
prevalence intensity of infection, 950
assay,
Ctenocephalides felis, 571
development, 571
larvae, 571
standards, 571
attractant,
Blattella germanica, 427
German cockroach, 427
methyl neoalkanamide, 427
repellent, 427
attractants,
canopy trap, 984
Haematopota pluvialis, 984
Hybomitra expollicata, 984
Morellia, 984
attraction,
feces, 957
stable fly, 957
Stomoxys calcitrans, 957
triple cage olfactometer, 957
Australia, 146, 153, 380, 736, 846
Aedes aegypti, 846
Aedes notoscriptus, 380
Anopheles farauti, 153
arbovirus, 846
Barmah Forest virus, 380
biochemical taxonomy, 736
Culex (Culex) sitiens subgroup, 736
Culicidae, 736
dispersal, 380
malaria, 153
mark-release-recapture, 380
Mesocyclops, 846
mosquito, 846
mosquitoes, 146
Papua New Guinea, 736
Ross River virus, 146, 380
seasonal abundance, 153
sibling species, 153
subterranean, 846
vector competence, 146
autumn–winter,
abundance variation, 16
Aedes albifasciatus, 16
gonotrophic status, 16
avian serology,
California, 507
St. Louis encephalitis virus, 507
surveillance, 507
western equine encephalomyelitis virus, 507

wetlands, 507
avoidance,
arrestant, 472
deer, 472
Ixodes scapularis, 472
kairomone, 472
ticks, 472

B

Bacillus sphaericus, 401, 534
bacterial insecticide, 401
Culex quinquefasciatus, 401, 534
insecticide resistance, 534
mosquitoes, 534
resistance, 401
resistance management, 401
bacteria, 246, 924
Anopheles gambiae, 246
Anopheles stephensi, 246
antibiotics, 246
artificial medium, 924
house fly, 924
intestinal tract, 924
larval development, 924
bacterial insecticide,
Bacillus sphaericus, 401
Culex quinquefasciatus, 401
resistance, 401
resistance management, 401
Barmah Forest virus, 380, 660
Aedes aegypti, 660
Aedes notoscriptus, 380, 660
Australia, 380
Culex annulirostris, 660
Culex quinquefasciatus, 660
Culex sitiens, 660
dispersal, 380
mark-release-recapture, 380
Ross River virus, 380
barrier islands,
Amblyomma americanum, 276
coastal Georgia, 276
Ehrlichia chaffeensis, 276
human monocytic ehrlichiosis, 276
prevalence, 276
bat,
Indiana, 445
Myotis sodalis, 445
parasitic mite, 445
Paraspinturnix globosa, 445
Spinturnicidae, 445
bats,
Aedes aegypti, 965
antibodies, 965
dengue virus, 965
neutralizing, 965
behavior,
adaptations, 114
Dermacentor albipictus, 114
host-finding, 114
larvae, 114
transmission, 114
bias,
abundance, 299
accuracy, 299
Haematobia irritans, 299
precision, 299
sampling, 299
bighorn sheep,
internal transcriber spacer region, 791
parasitism, 791
phylogeny, 791
Psoroptes, 791
biochemical taxonomy,
Australia, 736
Culex (Culex) sitiens subgroup, 736
Culicidae, 736
Papua New Guinea, 736
bioenhanced transmission,
enterohemorrhagic *Escherichia coli* O157: H7, 945
excretion, 945
house fly, 945
biological control,
insect picornaviruses, 271
Triatoma virus, 271
birds,
Borrelia burgdorferi, 909
Ixodes pacificus, 909
mice, 909
woodrats, 909
bite,
Aedes albopictus, 637
alightment, 637
mosquito attractant, 637
mosquito repellent, 637
proboscis amputation, 637
biting midge,
Culicoides sonorensis, 784
oral susceptibility, 784
vector competence, 784

vesicular stomatitis virus, 784
biting midges, 58, 65
aquatic ecology, 58
aquatic insects, 58
bluetongue, 65
Culicoides variipennis complex, 65
soil chemistry, 58
systematic, 65
vector distribution, 58
vectors, 65
Blattella germanica,
attractant, 427
German cockroach, 427
methyl neoalkanamide, 427
repellent, 427
blood feeding, 492, 743
Aedes aegypti, 492
DNA fingerprinting, 492
hematophagy, 743
hyaluronidase, 743
Phlebotomus papatasi, 743
salivary glands, 743
Simulium vittatum, 743
blood meal,
cat, 634
Ctenocephalides felis felis, 634
duration, 634
precocity, 634
blood-feeding,
antithrombin, 416
Hematobia irritans, 416
bluetongue,
biting midges, 65
Culicoides variipennis complex, 65
systematic, 65
vectors, 65
bluetongue virus,
Culicoides sonorensis, 675
seasonal abundance, 675
survivorship, 675
vectorial capacity, 675
body size,
Culex pipiens quinquefasciatus, 732
density, 732
life history traits, 732
body weight,
Dermacentor variabilis, 198
development, 198
Ixodes scapularis, 198
sex, 198
tick, 198
Borrelia,
diel activity, 9
litter, 9
season, 9
temperature gradient, 9
tick, 9
Borrelia burgdorferi, 102, 265, 766, 909
antimicrobial peptides, 265
birds, 909
Dermacentor variabilis, 265
detection, 102
Ixodes muris, 766
Ixodes pacificus, 909
Ixodes scapularis, 102
Lyme disease, 102
mice, 909
phagocytosis, 265
ribosomal RNA, 102
vector competence, 766
woodrats, 909
Brazil,
Amblyomma cajennense, 774
Amblyomma oblongoguttatum, 774
Amblyomma ovale, 774
dogs, 774
ticks, 774

C

California, 484, 507
avian serology, 507
drag sampling, 484
efficiency, 484
Ixodes pacificus, 484
nymphs, 484
St. Louis encephalitis virus, 507
surveillance, 507
western equine encephalomyelitis virus, 507
wetlands, 507
cannibalism,
life tables, 385
mortality, 385
survivorship, 385
Toxorhynchites rutilus, 385
canopy trap,
attractants, 984
Haematopota pluvialis, 984
Hybomitra expollicata, 984
Morellia, 984
capillary feeding,
Amblyomma maculatum, 761
Ixodid ticks, 761

production of aggregation-attachment pheromones, 761
Carcinops pumilio,
dispersal, 929
house fly, 929
cat,
blood meal, 634
Ctenocephalides felis felis, 634
duration, 634
precocity, 634
cat flea,
Ctenocephalides felis, 828
fecundity, 828
fertility, 828
multiple mating, 828
reproduction, 828
cat fleas,
Ctenocephalides felis, 503
ovisterilant larvicidal toxicity, 503
pyriproxyfen ingestion, 503
catalytic properties,
Dipetalogaster maximus, 689
flight muscle, 689
hexokinase, 689
cattle, 27, 210
chronobiology, 210
enzyme-linked immunosorbent assay, 27
experimental cycle, 210
Hypoderma bovis, 210
Hypoderma lineatum, 210
Hypoderma spp., 27
immunoresponse, 210
seroprevalence, 27
Spain, 27
warble flies, 27
cellular activity,
histology, 393
Hyalomma dromedarii, 393
permethrin, 393
CGA-255'728,
Ctenocephalides felis, 141
flea development, 141
insect growth regulator, 141
juvenile hormone mimic, 141
Chagas disease,
isoenzymes, 872
morphometrics, 872
population genetics, 872
random amplified polymorphic DNA, 872
Triatoma brasiliensis, 872
chromosomes,
hybridization, 29
laboratory rearing, 29
multiple mating, 29
Simulium damnosum complex, 29
chronobiology,
cattle, 210
experimental cycle, 210
Hypoderma bovis, 210
Hypoderma lineatum, 210
immunoresponse, 210
cibarial armature,
Anopheles flavirostris, 476
Anopheles minimus, 476
hybridization, 476
Indonesia, 476
circumsporozoite protein,
detection, 581
dipstick, 581
malaria, 581
mosquito, 581
classification,
endosymbiont, 340
mosquito, 340
polymerase chain reaction, 340
Wolbachia, 340
wsp, 340
climate,
interrelationship, 820
Ixodes didelphidis, 820
Ixodes loricatus, 820
small mammals, 820
coadaptation,
aedeagal filaments, 653
arms race, 653
Phlebotomine sand flies, 653
spermathecal ducts, 653
coastal Georgia,
Amblyomma americanum, 276
barrier islands, 276
Ehrlichia chaffeensis, 276
human monocytic ehrlichiosis, 276
prevalence, 276
Coelomomyces numularius,
Angola, 962
Anopheles squamosus, 962
new species, 962
COI gene,
Anopheles punctipennis, 754
mitochondrial DNA, 754

population structure, 754
colonization,
Greece, 346
Phlebotomus neglectus, 346
competition,
Aedes albopictus, 454
Aedes triseriatus, 454
invasion, 454
La Crosse virus, 454
oviposition, 454
concanavalin A,
acquired resistance, 885
Ixodes scapularis, 885
lipopolysaccharide, 885
lymphocytes, 885
salivary gland, 885
control, 121, 228
acaricides, 121
Amblyomma americanum, 121
Ctenocephalides felis, 228
ectoparasite, 228
imidacloprid, 228
Ixodes scapularis, 121
Lyme disease, 121
Coquillettidia perturbans,
Aedes albopictus, 701
Anopheles quadrimaculatus, 701
eastern equine encephalomyelitis virus, 701
survivorship, 701
coronavirus,
darkling beetle, 480
lesser mealworm, 480
litter beetle, 480
poult enteritis mortality syndrome, 480
turkey disease, 480
Costa Rica,
diversity, 216
Leishmaniasis, 216
Lutzomyia, 216
species composition, 216
cross-reactions,
Dermatophagoides farinae, 35
Dermatophagoides pteronyssinus, 35
house dust mites, 35
mite allergens, 35
mite growth, 35
cross-resistance,
insecticide resistance, 721
Pediculus capitis, 721
synergism, 721
cryopreservation,
adult, 968
primary screwworm, 968
pupa, 968
survival, 968
Ctenocephalides felis, 141, 228, 503, 571, 828
assay, 571
cat flea, 828
cat fleas, 503
CGA-255'728, 141
control, 228
development, 571
ectoparasite, 228
fecundity, 828
fertility, 828
flea development, 141
imidacloprid, 228
insect growth regulator, 141
juvenile hormone mimic, 141
larvae, 571
multiple mating, 828
ovisterilant larvicidal toxicity, 503
pyriproxyfen ingestion, 503
reproduction, 828
standards, 571
Ctenocephalides felis felis,
blood meal, 634
cat, 634
duration, 634
precocity, 634
Culex (Culex) sitiens subgroup,
Australia, 736
biochemical taxonomy, 736
Culicidae, 736
Papua New Guinea, 736
Culex annulirostris, 660, 797
Aedes aegypti, 660
Aedes notoscriptus, 660
Barmah Forest virus, 660
Culex quinquefasciatus, 660
Culex sitiens, 660
epidemiology, 797
Japanese encephalitis, 797
mosquito dispersal, 797
Murray Valley encephalitis, 797
Culex pipiens molestus, 726, 780
Anopheles stephensi, 726
Culex pipiens pallens, 726
Culex tritaeniorhynchus, 780

fatty acids, 726
Japanese encephalitis virus, 780
lipids, 726
susceptibility, 780
Taiwan, 780
yolk deposition, 726
Culex pipiens pallens,
Anopheles stephensi, 726
Culex pipiens molestus, 726
fatty acids, 726
lipids, 726
yolk deposition, 726
Culex pipiens quinquefasciatus,
body size, 732
density, 732
life history traits, 732
Culex quinquefasciatus, 172, 201, 401, 534, 660, 769
Aedes aegypti, 660
Aedes notoscriptus, 660
Aedes polynesiensis, 769
age-grading, 769
airborne action, 201
Anopheles culicifacies, 201
Bacillus sphaericus, 401, 534
bacterial insecticide, 401
Barmah Forest virus, 660
Culex annulirostris, 660
Culex sitiens, 660
excito-repellency action, 201
fluorescence, 769
gravid trap, 172
insecticide resistance, 534
insecticide-treated mosquito nets, 201
killing action, 201
mosquitoes, 534
pteridines, 769
resistance, 401
resistance management, 401
spectrofluorometry, 769
Tanzania, 172
Culex sitiens,
Aedes aegypti, 660
Aedes notoscriptus, 660
Barmah Forest virus, 660
Culex annulirostris, 660
Culex quinquefasciatus, 660
Culex tarsalis, 187, 194, 250
antibody, 250
Iowa, 194
longevity, 187
mosquito blood feeding, 250
multiplication, 194
spontaneous flight activity, 187
St. Louis encephalitis virus, 250
susceptibility, 194
transmission, 194
viremia, 250
western equine encephalomyelitis virus, 187, 194, 250
Culex tritaeniorhynchus,
Culex pipiens molestus, 780
Japanese encephalitis virus, 780
susceptibility, 780
Taiwan, 780
Culex vishnui subgroup,
identification, 554
polymerase chain reaction, 554
ribosomal DNA, 554
culicidae,
Aedes aegypti, 541
Culicidae, 736, 893
Australia, 736
biochemical taxonomy, 736
Culex (Culex) sitiens subgroup, 736
meconium, 893
metamorphosis, 893
mosquito, 893
Papua New Guinea, 736
peritrophic membrane, 893
culicidae,
resting behavior, 541
ultra-low volume insecticides, 541
Culicoides impunctatus,
antifeedant, 222
6-methylsalicylic acid, 222
repellent, 222
salicylic acid, 222
salicyluric acid, 222
Culicoides sonorensis, 675, 784
biting midge, 784
bluetongue virus, 675
oral susceptibility, 784
seasonal abundance, 675
survivorship, 675
vector competence, 784
vectorial capacity, 675
vesicular stomatitis virus, 784
Culicoides variipennis complex,
biting midges, 65
bluetongue, 65

systematic, 65
vectors, 65
Culiseta incidens,
Dirofilaria immitis, 295
heartworm, 295
Los Angeles County, 295
vectors, 295
Culiseta melanura,
arbovirus infection, 815
eastern equine encephalomyelitis virus, 815
mosquito mortality, 815
virulence, 815
culture,
eggs, 316
Forcipomyia (*Lasiohelea*) *townsvillensis*, 316
hatching, 316
oviposition, 316
cutaneous myiasis,
sarcophagid, 854
scanning electron microscopy, 854
cytoplasmic incompatibility,
Armigeres subalbatus, 53
Wolbachia, 53

D

darkling beetle,
coronavirus, 480
lesser mealworm, 480
litter beetle, 480
poult enteritis mortality syndrome, 480
turkey disease, 480
Dasypus novemcinctus,
Amblyomma auricularium, 977
armadillo, 977
Florida, 977
DDT,
Anopheles arabiensis, 286
Anopheles gambiae complex, 286
susceptibility, 286
synthetic pyrethroids, 286
deer,
arrestant, 472
avoidance, 472
Ixodes scapularis, 472
kairomone, 472
ticks, 472
deet,
Aedes communis, 919
AI3-37220, 919
piperidine, 919
repellents, 919
Simulium venustum, 919
degree-day,
Aedes albopictus, 40
Aedes triseriatus, 40
immature development, 40
dengue virus, 641, 965
Aedes aegypti, 641, 965
antibodies, 965
antibody effect, 641
anti-dengue serum, 641
bats, 965
neutralizing, 965
virus isolation methods, 641
density,
body size, 732
Culex pipiens quinquefasciatus, 732
life history traits, 732
Dermacentor albipictus,
adaptations, 114
behavior, 114
host-finding, 114
larvae, 114
transmission, 114
Dermacentor variabilis, 198, 265, 439
Amblyomma americanum, 439
Amblyomma maculatum, 439
antimicrobial peptides, 265
body weight, 198
Borrelia burgdorferi, 265
development, 198
Ixodes scapularis, 198, 439
phagocytosis, 265
sex, 198
tick, 198
tick ecology, 439
Dermatophagoides farinae,
cross-reactions, 35
Dermatophagoides pteronyssinus, 35
house dust mites, 35
mite allergens, 35
mite growth, 35
Dermatophagoides pteronyssinus,
cross-reactions, 35
Dermatophagoides farinae, 35
house dust mites, 35
mite allergens, 35
mite growth, 35
detection, 102, 581
Borrelia burgdorferi, 102

circumsporozoite protein, 581
dipstick, 581
Ixodes scapularis, 102
Lyme disease, 102
malaria, 581
mosquito, 581
ribosomal RNA, 102
development, 134, 198, 571
assay, 571
body weight, 198
Ctenocephalides felis, 571
Dermacentor variabilis, 198
Ixodes scapularis, 198
larvae, 571
Leishmania amazonensis, 134
Leishmania braziliensis, 134
Lutzomyia migonei, 134
sex, 198
standards, 571
tick, 198
diel activity,
Borrelia, 9
litter, 9
season, 9
temperature gradient, 9
tick, 9
Dipetalogaster maximus, 689, 938
catalytic properties, 689
fatty acid transport, 938
fatty acid-binding protein, 938
flight muscle, 689
hexokinase, 689
muscles, 938
triatominae, 938
dipstick,
circumsporozoite protein, 581
detection, 581
malaria, 581
mosquito, 581
Dirofilaria immiti,
Aedes sollicitans, 695
Aedes taeniorhynchus, 695
dog heartworm, 695
mosquitoes, 695
vector competence, 695
Dirofilaria immitis,
Culiseta incidens, 295
heartworm, 295
Los Angeles County, 295
vectors, 295
dispersal, 380, 929
Aedes notoscriptus, 380
Australia, 380
Barmah Forest virus, 380
Carcinops pumilio, 929
house fly, 929
mark-release-recapture, 380
Ross River virus, 380
distribution,
Forcipomyia taiwana, 205
population abundance, 205
rainfall, 205
temperature, 205
diversity,
Costa Rica, 216
Leishmaniasis, 216
Lutzomyia, 216
species composition, 216
DNA fingerprinting,
Aedes aegypti, 492
blood feeding, 492
dog heartworm,
Aedes sollicitans, 695
Aedes taeniorhynchus, 695
Dirofilaria immiti, 695
mosquitoes, 695
vector competence, 695
dogs,
Amblyomma cajennense, 774
Amblyomma oblongoguttatum, 774
Amblyomma ovale, 774
Brazil, 774
ticks, 774
drag sampling,
California, 484
efficiency, 484
Ixodes pacificus, 484
nymphs, 484
duration,
blood meal, 634
cat, 634
Ctenocephalides felis felis, 634
precocity, 634

E

eastern equine encephalomyelitis virus, 701, 815
Aedes albopictus, 701
Anopheles quadrimaculatus, 701
arbovirus infection, 815
Coquillettidia perturbans, 701
Culiseta melanura, 815
mosquito mortality, 815

survivorship, 701
virulence, 815
ecology,
Armigeres subalbatus, 108
Japanese encephalitis virus, 108
rice-free, 108
Taiwan, 108
ectoparasite,
control, 228
Ctenocephalides felis, 228
imidacloprid, 228
ectoparasites,
Guinea-Bissau, 973
hosts, 973
Ixodidae, 973
ticks, 973
efficiency,
California, 484
drag sampling, 484
Ixodes pacificus, 484
nymphs, 484
eggs,
culture, 316
Forcipomyia (*Lasiohelea*) *townsvillensis*, 316
hatching, 316
oviposition, 316
Ehrlichia chaffeensis, 276, 349, 595
Amblyomma americanum, 276, 349, 595
barrier islands, 276
coastal Georgia, 276
human monocytic ehrlichiosis, 276, 349, 595
indirect immunofluorescence assay, 595
Maryland, 349
polymerase chain reaction, 349, 595
prevalence, 276
Ehrlichia risticii,
aquatic insect host, 619
Potomac horse fever, 619
trematode vector, 619
embryonic development,
anomalies, 807
Hyalomma marginatum, 807
teratology, 807
endosymbiont,
classification, 340
mosquito, 340
polymerase chain reaction, 340
Wolbachia, 340
wsp, 340
enterohemorrhagic *Escherichia coli* O157: H7,
bioenhanced transmission, 945
excretion, 945
house fly, 945
environmental factors, 231, 461
Anopheles aquasalis, 231
Anopheles oswaldoi, 231
Haematobia irritans, 461
larval habitats, 231
malaria, 231
ovipositional behavior, 461
spatial temporal distribution, 231
enzyme-linked immunosorbent assay,
cattle, 27
Hypoderma spp., 27
seroprevalence, 27
Spain, 27
warble flies, 27
epidemic,
altitude, 664
Anopheles, 664
forecasting methods, 664
malaria, 664
sample size, 664
epidemiology,
Culex annulirostris, 797
Japanese encephalitis, 797
mosquito dispersal, 797
Murray Valley encephalitis, 797
excito-repellency action,
airborne action, 201
Anopheles culicifacies, 201
Culex quinquefasciatus, 201
insecticide-treated mosquito nets, 201
killing action, 201
excretion,
bioenhanced transmission, 945
enterohemorrhagic *Escherichia coli* O157: H7, 945
house fly, 945
experimental cycle,
cattle, 210
chronobiology, 210
Hypoderma bovis, 210
Hypoderma lineatum, 210
immunoresponse, 210

F

fatty acid transport,
Dipetalogaster maximus, 938
fatty acid-binding protein, 938
muscles, 938
triatominae, 938
fatty acid-binding protein,
Dipetalogaster maximus, 938
fatty acid transport, 938
muscles, 938
triatominae, 938
fatty acids,
Anopheles stephensi, 726
Culex pipiens molestus, 726
Culex pipiens pallens, 726
lipids, 726
yolk deposition, 726
feces,
attraction, 957
stable fly, 957
Stomoxys calcitrans, 957
triple cage olfactometer, 957
fecundity, 373, 828
cat flea, 828
Ctenocephalides felis, 828
feeding source, 373
fertility, 828
life cycle, 373
multiple mating, 828
reproduction, 828
Triatoma, 373
fecundity immunity to ticks,
feeding, 71
Rhipicephalus sanguineus, 71
tick performance, 71
fecundity midgut,
feeding, 601
Rhipicephalus sanguineus, 601
salivary gland immunity to ticks, 601
tick performance, 601
feeding, 71, 601
fecundity immunity to ticks, 71
fecundity midgut, 601
Rhipicephalus sanguineus, 71, 601
salivary gland immunity to ticks, 601
tick performance, 71, 601
feeding source,
fecundity, 373
life cycle, 373
Triatoma, 373
fertility,
cat flea, 828
Ctenocephalides felis, 828
fecundity, 828
multiple mating, 828
reproduction, 828
flea development,
CGA-255'728, 141
Ctenocephalides felis, 141
insect growth regulator, 141
juvenile hormone mimic, 141
fleas, 422, 864
juvenile hormone analog, 864
Oropsylla hirsuta, 864
prairie dogs, 864
pyriproxyfen, 864
quantitative competitive polymerase chain reaction, 422
Yersinia pestis, 422
flies,
Mexico, 489
Musca domestica, 489
Taenia solium, 489
vectors, 489
flight muscle,
catalytic properties, 689
Dipetalogaster maximus, 689
hexokinase, 689
Florida,
Amblyomma auricularium, 977
armadillo, 977
Dasypus novemcinctus, 977
fluorescence,
Aedes polynesiensis, 769
age-grading, 769
Culex quinquefasciatus, 769
pteridines, 769
spectrofluorometry, 769
Forcipomyia (*Lasiohelea*) *townsvillensis*,
culture, 316
eggs, 316
hatching, 316
oviposition, 316
Forcipomyia taiwana,
distribution, 205
population abundance, 205
rainfall, 205
temperature, 205
forecasting methods,
altitude, 664

Anopheles, 664
epidemic, 664
malaria, 664
sample size, 664

G

genetic clines,
Aedes abserratus, 902
Aedes hexodontus, 902
Aedes punctor, 902
allozymes, 902
larval morphology, 902
genetic diversity,
Androctonus, 787
ITS, 787
ribosomal DNA, 787
scorpions, 787
genetic variation,
isozymes, 325
Lutzomyia longipalpis, 325
sand flies, 325
Venezuela, 325
German cockroach,
attractant, 427
Blattella germanica, 427
methyl neoalkanamide, 427
repellent, 427
Glossina morsitans centralis,
Glossina morsitans morsitans, 897
Glossina swynnertoni, 897
hybridization asymmetry, 897
tsetse, 897
Glossina morsitans morsitans,
Glossina morsitans centralis, 897
Glossina swynnertoni, 897
hybridization asymmetry, 897
tsetse, 897
Glossina swynnertoni,
Glossina morsitans centralis, 897
Glossina morsitans morsitans, 897
hybridization asymmetry, 897
tsetse, 897
gonotrophic status,
abundance variation, 16
Aedes albifasciatus, 16
autumn–winter, 16
gravid trap,
Culex quinquefasciatus, 172
Tanzania, 172
Greece,
colonization, 346
Phlebotomus neglectus, 346
Guinea-Bissau,
ectoparasites, 973
hosts, 973
Ixodidae, 973
ticks, 973
gynandromorphism,
Amblyomma oblongoguttatum, 777
ticks, 777

H

habitat drying,
Aedes triseriatus, 950
Ascogregarina barretti, 950
prevalence intensity of infection, 950
habitat structure,
Aedes canadensis, 559
Aedes triseriatus, 559
La Crosse encephalitis virus, 559
mosquito ecology, 559
Haematobia irritans, 299, 461
abundance, 299
accuracy, 299
bias, 299
environmental factors, 461
ovipositional behavior, 461
precision, 299
sampling, 299
Haematopota pluvialis,
attractants, 984
canopy trap, 984
Hybomitra expollicata, 984
Morellia, 984
hatching,
culture, 316
eggs, 316
Forcipomyia (*Lasiohelea*) *townsvillensis*, 316
oviposition, 316
head lice,
nicotine, 170
Pediculus humanus capitis, 170
heartworm,
Culiseta incidens, 295
Dirofilaria immitis, 295
Los Angeles County, 295
vectors, 295
Hematobia irritans,
antithrombin, 416
blood-feeding, 416
hematophagy,
blood feeding, 743

hyaluronidase, 743
Phlebotomus papatasi, 743
salivary glands, 743
Simulium vittatum, 743
heteroduplex analysis,
mitochondrial genes, 547
Onchocerca volvulus, 547
phylogeny, 547
Simulium damnosum, 547
Sudan, 547
heterologous promoters,
luciferase expression, 802
Lutzomyia, 802
Phlebotomus, 802
sand fly cells, 802
hexokinase,
catalytic properties, 689
Dipetalogaster maximus, 689
flight muscle, 689
histology, 89, 393
Aedes aegypti, 89
cellular activity, 393
Hyalomma dromedarii, 393
multiple blood feeding, 89
permethrin, 393
Puerto Rico, 89
Thailand, 89
host,
Manis, 640
parasite, 640
Sus, 640
Thailand, 640
tick, 640
host-finding,
adaptations, 114
behavior, 114
Dermacentor albipictus, 114
larvae, 114
transmission, 114
host-location behavior,
human odors, 308
stable fly, 308
Stomoxys calcitrans, 308
triple cage olfactometer, 308
hosts, 714, 973
ectoparasites, 973
Guinea-Bissau, 973
Ixodes (*Ixodes*) *loricatus*, 714
Ixodidae, 973
life cycle, 714
ticks, 973
house dust mites,
cross-reactions, 35
Dermatophagoides farinae, 35
Dermatophagoides pteronyssinus, 35
mite allergens, 35
mite growth, 35
house finch,
antibody, 259
infection, 259
St. Louis encephalitis virus, 259
viremia, 259
western equine encephalomyelitis virus, 259
house fly, 924, 929, 945
artificial medium, 924
bacteria, 924
bioenhanced transmission, 945
Carcinops pumilio, 929
dispersal, 929
enterohemorrhagic *Escherichia coli* O157: H7, 945
excretion, 945
intestinal tract, 924
larval development, 924
human monocytic ehrlichiosis, 276, 349, 595
Amblyomma americanum, 276, 349, 595
barrier islands, 276
coastal Georgia, 276
Ehrlichia chaffeensis, 276, 349, 595
indirect immunofluorescence assay, 595
Maryland, 349
polymerase chain reaction, 349, 595
prevalence, 276
human odors,
host-location behavior, 308
stable fly, 308
Stomoxys calcitrans, 308
triple cage olfactometer, 308
Hyalomma dromedarii,
cellular activity, 393
histology, 393
permethrin, 393
Hyalomma marginatum,
anomalies, 807
embryonic development, 807
teratology, 807
hyaluronidase,
blood feeding, 743

hematophagy, 743
Phlebotomus papatasi, 743
salivary glands, 743
Simulium vittatum, 743
Hybomitra expollicata,
attractants, 984
canopy trap, 984
Haematopota pluvialis, 984
Morellia, 984
hybridization, 29, 476
Anopheles flavirostris, 476
Anopheles minimus, 476
chromosomes, 29
cibarial armature, 476
Indonesia, 476
laboratory rearing, 29
multiple mating, 29
Simulium damnosum complex, 29
hybridization asymmetry,
Glossina morsitans centralis, 897
Glossina morsitans morsitans, 897
Glossina swynnertoni, 897
tsetse, 897
Hypoderma bovis,
cattle, 210
chronobiology, 210
experimental cycle, 210
Hypoderma lineatum, 210
immunoresponse, 210
Hypoderma lineatum,
cattle, 210
chronobiology, 210
experimental cycle, 210
Hypoderma bovis, 210
immunoresponse, 210
Hypoderma spp.,
cattle, 27
enzyme-linked immunosorbent assay, 27
seroprevalence, 27
Spain, 27
warble flies, 27

I

identification,
Culex vishnui subgroup, 554
polymerase chain reaction, 554
ribosomal DNA, 554
imidacloprid,
control, 228
Ctenocephalides felis, 228
ectoparasite, 228
immature development,
Aedes albopictus, 40
Aedes triseriatus, 40
degree-day, 40
immunoresponse,
cattle, 210
chronobiology, 210
experimental cycle, 210
Hypoderma bovis, 210
Hypoderma lineatum, 210
Indian sand flies,
antenna, 575
larvae, 575
mouthparts, 575
Phlebotomus, 575
Sergentomyia, 575
Indiana,
bat, 445
Myotis sodalis, 445
parasitic mite, 445
Paraspinturnix globosa, 445
Spinturnicidae, 445
indirect immunofluorescence assay,
Amblyomma americanum, 595
Ehrlichia chaffeensis, 595
human monocytic ehrlichiosis, 595
polymerase chain reaction, 595
Indonesia,
Anopheles flavirostris, 476
Anopheles minimus, 476
cibarial armature, 476
hybridization, 476
infection,
antibody, 259
house finch, 259
St. Louis encephalitis virus, 259
viremia, 259
western equine encephalomyelitis virus, 259
insect container communities,
Aedes aegypti, 364
Aedes albopictus, 364
invertebrate carcasses, 364
replacement series, 364
insect growth regulator,
CGA-255'728, 141
Ctenocephalides felis, 141
flea development, 141
juvenile hormone mimic, 141
insect picornaviruses,
biological control, 271

Triatoma virus, 271
insecticide resistance, 534, 721
Bacillus sphaericus, 534
cross-resistance, 721
Culex quinquefasciatus, 534
mosquitoes, 534
Pediculus capitis, 721
synergism, 721
insecticide-treated mosquito nets,
airborne action, 201
Anopheles culicifacies, 201
Culex quinquefasciatus, 201
excito-repellency action, 201
killing action, 201
internal transcriber spacer region,
bighorn sheep, 791
parasitism, 791
phylogeny, 791
Psoroptes, 791
interrelationship,
climate, 820
Ixodes didelphidis, 820
Ixodes loricatus, 820
small mammals, 820
intestinal tract,
artificial medium, 924
bacteria, 924
house fly, 924
larval development, 924
invasion,
Aedes albopictus, 454
Aedes triseriatus, 454
competition, 454
La Crosse virus, 454
oviposition, 454
invertebrate carcasses,
Aedes aegypti, 364
Aedes albopictus, 364
insect container communities, 364
replacement series, 364
Iowa,
Culex tarsalis, 194
multiplication, 194
susceptibility, 194
transmission, 194
western equine encephalomyelitis virus, 194
isoenzymes, 289, 872
Chagas disease, 872
morphometrics, 872
Phlebotomus brevis, 289
Phlebotomus simici, 289
population genetics, 872
random amplified polymorphic DNA, 872
sand flies, 289
scanning electron microscopy, 289
Triatoma brasiliensis, 872
isozymes,
genetic variation, 325
Lutzomyia longipalpis, 325
sand flies, 325
Venezuela, 325
ITS,
Androctonus, 787
genetic diversity, 787
ribosomal DNA, 787
scorpions, 787
ITS2,
Anopheles dispar, 933
Anopheles greeni, 933
Anopheles maculatus, 933
Philippines, 933
polymerase chain reaction-restriction fragment-length polymorphism, 933
ivermectin,
Ixodes scapularis, 126
Lyme disease, 126
tick control, 126
white-tailed deer, 126
Ixodes (*Ixodes*) *loricatus*,
hosts, 714
life cycle, 714
Ixodes didelphidis,
climate, 820
interrelationship, 820
Ixodes loricatus, 820
small mammals, 820
Ixodes loricatus,
climate, 820
interrelationship, 820
Ixodes didelphidis, 820
small mammals, 820
Ixodes muris,
Borrelia burgdorferi, 766
vector competence, 766
Ixodes pacificus, 484, 909
birds, 909
Borrelia burgdorferi, 909
California, 484
drag sampling, 484

efficiency, 484
mice, 909
nymphs, 484
woodrats, 909
Ixodes ricinus,
rickettsia, 707
spotted fever group, 707
*Ixodes ricinus*tick,
methodology, 335
sampling, 335
Ixodes scapularis, 102, 121, 126, 198, 357, 408, 439, 472, 645, 885
acaracide, 645
acaricides, 121
acquired resistance, 885
Amblyomma americanum, 121, 439
Amblyomma maculatum, 439
Amblyomma variegatum, 645
arrestant, 472
avoidance, 472
body weight, 198
Borrelia burgdorferi, 102
concanavalin A, 885
control, 121
deer, 472
Dermacentor variabilis, 198, 439
detection, 102
development, 198
ivermectin, 126
Ixodiphagus hookeri, 645
kairomone, 472
lipopolysaccharide, 885
Lyme disease, 102, 121, 126, 357, 408
lymphocytes, 885
parasitoid augmentation, 645
population dynamics, 408
population size, 357
ribosomal RNA, 102
salivary gland, 885
sampling efficiency, 357
sex, 198
tick, 198
tick control, 126
tick density, 357
tick ecology, 439
ticks, 472
weather, 408
white-footed mice, 408
white-tailed deer, 126
Ixodid ticks,
Amblyomma maculatum, 761
capillary feeding, 761
production of aggregation-attachment pheromones, 761
Ixodidae,
ectoparasites, 973
Guinea-Bissau, 973
hosts, 973
ticks, 973
Ixodiphagus hookeri,
acaracide, 645
Amblyomma variegatum, 645
Ixodes scapularis, 645
parasitoid augmentation, 645

J

Japanese encephalitis,
Culex annulirostris, 797
epidemiology, 797
mosquito dispersal, 797
Murray Valley encephalitis, 797
Japanese encephalitis virus, 108, 780
Armigeres subalbatus, 108
Culex pipiens molestus, 780
Culex tritaeniorhynchus, 780
ecology, 108
rice-free, 108
susceptibility, 780
Taiwan, 108, 780
juvenile hormone analog, 612, 864
fleas, 864
nontarget arthropods, 612
Oropsylla hirsuta, 864
prairie dogs, 864
pyriproxyfen, 612, 864
juvenile hormone mimic,
CGA-255'728, 141
Ctenocephalides felis, 141
flea development, 141
insect growth regulator, 141

K

kairomone,
arrestant, 472
avoidance, 472
deer, 472
Ixodes scapularis, 472
ticks, 472
killing action,
airborne action, 201
Anopheles culicifacies, 201
Culex quinquefasciatus, 201

excito-repellency action, 201
insecticide-treated mosquito nets, 201

L

La Crosse encephalitis virus,
Aedes canadensis, 559
Aedes triseriatus, 559
habitat structure, 559
mosquito ecology, 559
La Crosse virus,
Aedes albopictus, 454
Aedes triseriatus, 454
competition, 454
invasion, 454
oviposition, 454
laboratory rearing,
chromosomes, 29
hybridization, 29
multiple mating, 29
Simulium damnosum complex, 29
larvae, 114, 571, 575
adaptations, 114
antenna, 575
assay, 571
behavior, 114
Ctenocephalides felis, 571
Dermacentor albipictus, 114
development, 571
host-finding, 114
Indian sand flies, 575
mouthparts, 575
Phlebotomus, 575
Sergentomyia, 575
standards, 571
transmission, 114
larval development, 281, 924
Aedes albopictus, 281
artificial medium, 924
bacteria, 924
house fly, 924
intestinal tract, 924
metallic copper, 281
mosquito control, 281
larval habitats,
Anopheles aquasalis, 231
Anopheles oswaldoi, 231
environmental factors, 231
malaria, 231
spatial temporal distribution, 231
larval morphology,
Aedes abserratus, 902
Aedes hexodontus, 902
Aedes punctor, 902
allozymes, 902
genetic clines, 902
Leishmania amazonensis,
development, 134
Leishmania braziliensis, 134
Lutzomyia migonei, 134
Leishmania braziliensis,
development, 134
Leishmania amazonensis, 134
Lutzomyia migonei, 134
Leishmaniasis,
Costa Rica, 216
diversity, 216
Lutzomyia, 216
species composition, 216
leishmaniasis,
Lutzomyia spp., 162
Rio Grande do Norte, 162
sand flies, 162
lesser mealworm,
coronavirus, 480
darkling beetle, 480
litter beetle, 480
poult enteritis mortality syndrome, 480
turkey disease, 480
life cycle, 373, 714
fecundity, 373
feeding source, 373
hosts, 714
Ixodes (*Ixodes*) *loricatus*, 714
Triatoma, 373
life history traits,
body size, 732
Culex pipiens quinquefasciatus, 732
density, 732
life tables,
cannibalism, 385
mortality, 385
survivorship, 385
Toxorhynchites rutilus, 385
lipids,
Anopheles stephensi, 726
Culex pipiens molestus, 726
Culex pipiens pallens, 726
fatty acids, 726
yolk deposition, 726
lipopolysaccharide,
acquired resistance, 885

concanavalin A, 885
Ixodes scapularis, 885
lymphocytes, 885
salivary gland, 885
litter,
Borrelia, 9
diel activity, 9
season, 9
temperature gradient, 9
tick, 9
litter beetle,
coronavirus, 480
darkling beetle, 480
lesser mealworm, 480
poult enteritis mortality syndrome, 480
turkey disease, 480
lone star tick, 588, 878
acaricide, 588, 878
Amblyomma americanum, 588, 878
amitraz, 588, 878
self-treatment, 588, 878
longevity,
Culex tarsalis, 187
spontaneous flight activity, 187
western equine encephalomyelitis virus, 187
Los Angeles County,
Culiseta incidens, 295
Dirofilaria immitis, 295
heartworm, 295
vectors, 295
luciferase expression,
heterologous promoters, 802
Lutzomyia, 802
Phlebotomus, 802
sand fly cells, 802
Lutzomyia, 216, 860
Costa Rica, 216
diversity, 216
heterologous promoters, 802
Leishmaniasis, 216
luciferase expression, 802
Phlebotomus, 802
sand fly, 860
sand fly cells, 802
sensilla coeloconica, 860
species composition, 216
ultrastructure, 860
Lutzomyia longipalpis,
genetic variation, 325
isozymes, 325
sand flies, 325
Venezuela, 325
Lutzomyia migonei, 134, 748
development, 134
Leishmania amazonensis, 134
Leishmania braziliensis, 134
mouthparts, 748
sandflies, 748
Lutzomyia spp.,
leishmaniasis, 162
Rio Grande do Norte, 162
sand flies, 162
Lyme disease, 102, 121, 126, 357, 408
acaricides, 121
Amblyomma americanum, 121
Borrelia burgdorferi, 102
control, 121
detection, 102
ivermectin, 126
Ixodes scapularis, 102, 121, 126, 357, 408
population dynamics, 408
population size, 357
ribosomal RNA, 102
sampling efficiency, 357
tick control, 126
tick density, 357
weather, 408
white-footed mice, 408
white-tailed deer, 126
lymphocytes,
acquired resistance, 885
concanavalin A, 885
Ixodes scapularis, 885
lipopolysaccharide, 885
salivary gland, 885

M

malaria, 153, 231, 581, 664
altitude, 664
Anopheles, 664
Anopheles aquasalis, 231
Anopheles farauti, 153
Anopheles oswaldoi, 231
Australia, 153
circumsporozoite protein, 581
detection, 581
dipstick, 581
environmental factors, 231
epidemic, 664
forecasting methods, 664

larval habitats, 231
mosquito, 581
sample size, 664
seasonal abundance, 153
sibling species, 153
spatial temporal distribution, 231
Manis,
host, 640
parasite, 640
Sus, 640
Thailand, 640
tick, 640
mark-release-recapture,
Aedes notoscriptus, 380
Australia, 380
Barmah Forest virus, 380
dispersal, 380
Ross River virus, 380
Maryland,
Amblyomma americanum, 349
Ehrlichia chaffeensis, 349
human monocytic ehrlichiosis, 349
polymerase chain reaction, 349
meconium,
Culicidae, 893
metamorphosis, 893
mosquito, 893
peritrophic membrane, 893
Mesocyclops,
Aedes aegypti, 846
arbovirus, 846
Australia, 846
mosquito, 846
subterranean, 846
metallic copper,
Aedes albopictus, 281
larval development, 281
mosquito control, 281
metamorphosis,
Culicidae, 893
meconium, 893
mosquito, 893
peritrophic membrane, 893
methodology,
*Ixodes ricinus*tick, 335
sampling, 335
methyl neoalkanamide,
attractant, 427
Blattella germanica, 427
German cockroach, 427
repellent, 427
Mexico,
flies, 489
Musca domestica, 489
Taenia solium, 489
vectors, 489
mice,
birds, 909
Borrelia burgdorferi, 909
Ixodes pacificus, 909
woodrats, 909
mite allergens,
cross-reactions, 35
Dermatophagoides farinae, 35
Dermatophagoides pteronyssinus, 35
house dust mites, 35
mite growth, 35
mite growth,
cross-reactions, 35
Dermatophagoides farinae, 35
Dermatophagoides pteronyssinus, 35
house dust mites, 35
mite allergens, 35
mitochondrial DNA,
Anopheles punctipennis, 754
COI gene, 754
population structure, 754
mitochondrial genes,
heteroduplex analysis, 547
Onchocerca volvulus, 547
phylogeny, 547
Simulium damnosum, 547
Sudan, 547
Monoclonal antibody,
Anopheles gambiae, 528
phage display, 528
scFv, 528
screening, 528
transmission blocking, 528
Morellia,
attractants, 984
canopy trap, 984
Haematopota pluvialis, 984
Hybomitra expollicata, 984
morphology,
Oestrus ovis, 239
parasite, 239
puparium, 239
sheep nose bot, 239
morphometrics,
Chagas disease, 872
isoenzymes, 872

population genetics, 872
random amplified polymorphic DNA, 872
Triatoma brasiliensis, 872
mortality,
cannibalism, 385
life tables, 385
survivorship, 385
Toxorhynchites rutilus, 385
mosquito, 340, 581, 846, 893
Aedes aegypti, 846
arbovirus, 846
Australia, 846
circumsporozoite protein, 581
classification, 340
Culicidae, 893
detection, 581
dipstick, 581
endosymbiont, 340
malaria, 581
meconium, 893
Mesocyclops, 846
metamorphosis, 893
peritrophic membrane, 893
polymerase chain reaction, 340
subterranean, 846
Wolbachia, 340
wsp, 340
mosquito attractant,
Aedes albopictus, 637
alightment, 637
bite, 637
mosquito repellent, 637
proboscis amputation, 637
mosquito blood feeding,
antibody, 250
Culex tarsalis, 250
St. Louis encephalitis virus, 250
viremia, 250
western equine encephalomyelitis virus, 250
mosquito control,
Aedes albopictus, 281
larval development, 281
metallic copper, 281
mosquito dispersal,
Culex annulirostris, 797
epidemiology, 797
Japanese encephalitis, 797
Murray Valley encephalitis, 797
mosquito ecology,
Aedes canadensis, 559
Aedes triseriatus, 559
habitat structure, 559
La Crosse encephalitis virus, 559
mosquito mortality,
arbovirus infection, 815
Culiseta melanura, 815
eastern equine encephalomyelitis virus, 815
virulence, 815
mosquito repellent,
Aedes albopictus, 637
alightment, 637
bite, 637
mosquito attractant, 637
proboscis amputation, 637
mosquito-borne encephalitis,
arboviral transmission, 626
St. Louis encephalitis virus, 626
vector infection rates, 626
mosquitoes, 146, 467, 534, 695, 835
Aedes sollicitans, 695
Aedes taeniorhynchus, 695
Australia, 146
Bacillus sphaericus, 534
Culex quinquefasciatus, 534
Dirofilaria immiti, 695
dog heartworm, 695
insecticide resistance, 534
Peru, 835
reverse transcription-polymerase chain reaction, 467
Rift Valley fever virus, 467
Ross River virus, 146
transmission, 835
vector competence, 146, 695, 835
Venezuelan equine encephalomyelitis virus, 835
mouthparts, 575, 748
antenna, 575
Indian sand flies, 575
larvae, 575
Lutzomyia migonei, 748
Phlebotomus, 575
sandflies, 748
Sergentomyia, 575
multiple blood feeding,
Aedes aegypti, 89
histology, 89
Puerto Rico, 89

Thailand, 89
multiple mating, 29, 828
cat flea, 828
chromosomes, 29
Ctenocephalides felis, 828
fecundity, 828
fertility, 828
hybridization, 29
laboratory rearing, 29
reproduction, 828
Simulium damnosum complex, 29
multiplication,
Culex tarsalis, 194
Iowa, 194
susceptibility, 194
transmission, 194
western equine encephalomyelitis virus, 194
Murray Valley encephalitis,
Culex annulirostris, 797
epidemiology, 797
Japanese encephalitis, 797
mosquito dispersal, 797
Musca domestica,
flies, 489
Mexico, 489
Taenia solium, 489
vectors, 489
muscles,
Dipetalogaster maximus, 938
fatty acid transport, 938
fatty acid-binding protein, 938
triatominae, 938
Myotis sodalis,
bat, 445
Indiana, 445
parasitic mite, 445
Paraspinturnix globosa, 445
Spinturnicidae, 445

N

N,N-diethyl-3-methylbenzamide,
Anopheles stephensi, 177
1-[3-cyclohexen-1-ylcarbonyl] piperidine, 177
1-[3-cyclohexen-1-ylcarbonyl]-2-methylpiperidine, 177
new module, 177
neutralizing,
Aedes aegypti, 965
antibodies, 965
bats, 965
dengue virus, 965
new module,
1-[3-cyclohexen-1-ylcarbonyl] piperidine, 177
1-[3-cyclohexen-1-ylcarbonyl]-2-methylpiperidine, 177
Anopheles stephensi, 177
N,N-diethyl-3-methylbenzamide, 177
new species,
Angola, 962
Anopheles squamosus, 962
Coelomomyces numularius, 962
nicotine,
head lice, 170
Pediculus humanus capitis, 170
nonindigenous arthropods,
arthropod pest introductions, 1
arthropod-borne diseases, 1
nontarget arthropods,
juvenile hormone analog, 612
pyriproxyfen, 612
nymphs,
California, 484
drag sampling, 484
efficiency, 484
Ixodes pacificus, 484

O

Oestrus ovis,
morphology, 239
parasite, 239
puparium, 239
sheep nose bot, 239
Onchocerca volvulus,
heteroduplex analysis, 547
mitochondrial genes, 547
phylogeny, 547
Simulium damnosum, 547
Sudan, 547
oral susceptibility,
biting midge, 784
Culicoides sonorensis, 784
vector competence, 784
vesicular stomatitis virus, 784
Oropsylla hirsuta,
fleas, 864
juvenile hormone analog, 864
prairie dogs, 864
pyriproxyfen, 864
oviposition, 316, 454
Aedes albopictus, 454

Aedes triseriatus, 454
competition, 454
culture, 316
eggs, 316
Forcipomyia (*Lasiohelea*) *townsvillensis*, 316
hatching, 316
invasion, 454
La Crosse virus, 454
ovipositional behavior,
environmental factors, 461
Haematobia irritans, 461
ovisterilant larvicidal toxicity,
cat fleas, 503
Ctenocephalides felis, 503
pyriproxyfen ingestion, 503

P

Pantanal region,
ticks, 979
wild animals, 979
Papua New Guinea, 736, 840
Anopheles clowi, 840
Anopheles punctulatus group, 840
Australia, 736
biochemical taxonomy, 736
Culex (Culex) sitiens subgroup, 736
Culicidae, 736
parasite, 239, 640
host, 640
Manis, 640
morphology, 239
Oestrus ovis, 239
puparium, 239
sheep nose bot, 239
Sus, 640
Thailand, 640
tick, 640
parasitic mite,
bat, 445
Indiana, 445
Myotis sodalis, 445
Paraspinturnix globosa, 445
Spinturnicidae, 445
parasitism,
bighorn sheep, 791
internal transcriber spacer region, 791
phylogeny, 791
Psoroptes, 791
parasitoid augmentation,
acaracide, 645
Amblyomma variegatum, 645
Ixodes scapularis, 645
Ixodiphagus hookeri, 645
Paraspinturnix globosa,
bat, 445
Indiana, 445
Myotis sodalis, 445
parasitic mite, 445
Spinturnicidae, 445
Pediculus capitis,
cross-resistance, 721
insecticide resistance, 721
synergism, 721
Pediculus humanus capitis,
head lice, 170
nicotine, 170
peritrophic membrane,
Culicidae, 893
meconium, 893
metamorphosis, 893
mosquito, 893
permethrin,
cellular activity, 393
histology, 393
Hyalomma dromedarii, 393
Peru,
mosquitoes, 835
transmission, 835
vector competence, 835
Venezuelan equine encephalomyelitis virus, 835
phage display,
Anopheles gambiae, 528
Monoclonal antibody, 528
scFv, 528
screening, 528
transmission blocking, 528
phagocytosis,
antimicrobial peptides, 265
Borrelia burgdorferi, 265
Dermacentor variabilis, 265
phenotypes,
Anopheles albimanus, 331
Plasmodium vivax, 331
susceptibility, 331
Philippines,
Anopheles dispar, 933
Anopheles greeni, 933
Anopheles maculatus, 933
ITS2, 933
polymerase chain reaction-

restriction fragment-length polymorphism, 933
Phlebotomine sand flies,
aedeagal filaments, 653
arms race, 653
coadaptation, 653
spermathecal ducts, 653
Phlebotomus,
antenna, 575
heterologous promoters, 802
Indian sand flies, 575
larvae, 575
luciferase expression, 802
Lutzomyia, 802
mouthparts, 575
sand fly cells, 802
Sergentomyia, 575
Phlebotomus brevis,
isoenzymes, 289
Phlebotomus simici, 289
sand flies, 289
scanning electron microscopy, 289
Phlebotomus duboscqi,
Phlebotomus papatasi, 435
Phlebotomus sergenti, 435
Rift Valley fever virus, 435
Sergentomyia schwetzi, 435
vector competency, 435
Phlebotomus neglectus,
colonization, 346
Greece, 346
Phlebotomus papatasi,
blood feeding, 743
hematophagy, 743
hyaluronidase, 743
Phlebotomus papatasi, 319, 435
Phlebotomus duboscqi, 435
Phlebotomus sergenti, 435
photosynthesis, 319
plant feeding, 319
Rift Valley fever virus, 435
Phlebotomus papatasi,
salivary glands, 743
Phlebotomus papatasi,
Sergentomyia schwetzi, 435
Phlebotomus papatasi,
Simulium vittatum, 743
Phlebotomus papatasi, 319, 435
starch grains, 319
sugars, 319
vector competency, 435
Phlebotomus sergenti,
Phlebotomus duboscqi, 435
Phlebotomus papatasi, 435
Rift Valley fever virus, 435
Sergentomyia schwetzi, 435
vector competency, 435
Phlebotomus simici,
isoenzymes, 289
Phlebotomus brevis, 289
sand flies, 289
scanning electron microscopy, 289
photosynthesis,
Phlebotomus papatasi, 319
plant feeding, 319
starch grains, 319
sugars, 319
phylogeny, 547, 791
bighorn sheep, 791
heteroduplex analysis, 547
internal transcriber spacer region, 791
mitochondrial genes, 547
Onchocerca volvulus, 547
parasitism, 791
Psoroptes, 791
Simulium damnosum, 547
Sudan, 547
piperidine,
Aedes communis, 919
AI3-37220, 919
deet, 919
repellents, 919
Simulium venustum, 919
plant feeding,
Phlebotomus papatasi, 319
photosynthesis, 319
starch grains, 319
sugars, 319
Plasmodium vivax,
Anopheles albimanus, 331
phenotypes, 331
susceptibility, 331
polymerase chain reaction, 340, 349, 554, 595
Amblyomma americanum, 349, 595
classification, 340
Culex vishnui subgroup, 554
Ehrlichia chaffeensis, 349, 595
endosymbiont, 340
human monocytic ehrlichiosis, 349, 595

identification, 554
indirect immunofluorescence assay, 595
Maryland, 349
mosquito, 340
ribosomal DNA, 554
Wolbachia, 340
wsp, 340
polymerase chain reaction-restriction fragment-length polymorphism,
Anopheles dispar, 933
Anopheles greeni, 933
Anopheles maculatus, 933
ITS2, 933
Philippines, 933
population abundance,
distribution, 205
Forcipomyia taiwana, 205
rainfall, 205
temperature, 205
population dynamics, 21, 77, 408
Aedes aegypti, 77
Aedes albifasciatus, 21
Argentina, 21
Ixodes scapularis, 408
Lyme disease, 408
Puerto Rico, 77
Thailand, 77
weather, 408
white-footed mice, 408
population genetics,
Chagas disease, 872
isoenzymes, 872
morphometrics, 872
random amplified polymorphic DNA, 872
Triatoma brasiliensis, 872
population size,
Ixodes scapularis, 357
Lyme disease, 357
sampling efficiency, 357
tick density, 357
population structure,
Anopheles punctipennis, 754
COI gene, 754
mitochondrial DNA, 754
Potomac horse fever,
aquatic insect host, 619
Ehrlichia risticii, 619
trematode vector, 619
poult enteritis mortality syndrome,
coronavirus, 480
darkling beetle, 480
lesser mealworm, 480
litter beetle, 480
turkey disease, 480
prairie dogs,
fleas, 864
juvenile hormone analog, 864
Oropsylla hirsuta, 864
pyriproxyfen, 864
precision,
abundance, 299
accuracy, 299
bias, 299
Haematobia irritans, 299
sampling, 299
precocity,
blood meal, 634
cat, 634
Ctenocephalides felis felis, 634
duration, 634
prevalence,
Amblyomma americanum, 276
barrier islands, 276
coastal Georgia, 276
Ehrlichia chaffeensis, 276
human monocytic ehrlichiosis, 276
prevalence intensity of infection,
Aedes triseriatus, 950
Ascogregarina barretti, 950
habitat drying, 950
primary screwworm,
adult, 968
cryopreservation, 968
pupa, 968
survival, 968
proboscis amputation,
Aedes albopictus, 637
alightment, 637
bite, 637
mosquito attractant, 637
mosquito repellent, 637
production of aggregation-attachment pheromones,
Amblyomma maculatum, 761
capillary feeding, 761
Ixodid ticks, 761
Psoroptes,
bighorn sheep, 791
internal transcriber spacer region, 791

parasitism, 791
phylogeny, 791
pteridines,
Aedes polynesiensis, 769
age-grading, 769
Culex quinquefasciatus, 769
fluorescence, 769
spectrofluorometry, 769
Puerto Rico, 77, 89
Aedes aegypti, 77, 89
histology, 89
multiple blood feeding, 89
population dynamics, 77
Thailand, 77, 89
pupa,
adult, 968
cryopreservation, 968
primary screwworm, 968
survival, 968
puparium,
morphology, 239
Oestrus ovis, 239
parasite, 239
sheep nose bot, 239
pyriproxyfen, 612, 864
fleas, 864
juvenile hormone analog, 612, 864
nontarget arthropods, 612
Oropsylla hirsuta, 864
prairie dogs, 864
pyriproxyfen ingestion,
cat fleas, 503
Ctenocephalides felis, 503
ovisterilant larvicidal toxicity, 503

Q

quantitative competitive polymerase chain reaction,
fleas, 422
Yersinia pestis, 422

R

rainfall,
distribution, 205
Forcipomyia taiwana, 205
population abundance, 205
temperature, 205
random amplified polymorphic DNA,
Chagas disease, 872
isoenzymes, 872
morphometrics, 872
population genetics, 872
Triatoma brasiliensis, 872
repellent, 222, 427
antifeedant, 222
attractant, 427
Blattella germanica, 427
Culicoides impunctatus, 222
German cockroach, 427
6-methylsalicylic acid, 222
methyl neoalkanamide, 427
salicylic acid, 222
salicyluric acid, 222
repellents, 182, 919
Aedes communis, 919
AI3-37220, 919
Anopheles stephensi, 182
1-[3-cyclohexen-1-ylcarbonyl]-2-methylpiperidine, 182
1-[3-cyclohexen-1-ylcarbonyl] piperidine, 182
deet, 919
piperidine, 919
Simulium venustum, 919
stereoisomers, 182
replacement series,
Aedes aegypti, 364
Aedes albopictus, 364
insect container communities, 364
invertebrate carcasses, 364
reproduction,
cat flea, 828
Ctenocephalides felis, 828
fecundity, 828
fertility, 828
multiple mating, 828
resistance,
Bacillus sphaericus, 401
bacterial insecticide, 401
Culex quinquefasciatus, 401
resistance management, 401
resistance management,
Bacillus sphaericus, 401
bacterial insecticide, 401
Culex quinquefasciatus, 401
resistance, 401
resting behavior,
Aedes aegypti, 541
culicidae, 541
ultra-low volume insecticides, 541
reverse transcription-polymerase chain reaction,

mosquitoes, 467
Rift Valley fever virus, 467
Rhipicephalus sanguineus, 71, 601
fecundity immunity to ticks, 71
fecundity midgut, 601
feeding, 71, 601
salivary gland immunity to ticks, 601
tick performance, 71, 601
ribosomal DNA, 554, 787
Androctonus, 787
Culex vishnui subgroup, 554
genetic diversity, 787
identification, 554
ITS, 787
polymerase chain reaction, 554
scorpions, 787
ribosomal RNA,
Borrelia burgdorferi, 102
detection, 102
Ixodes scapularis, 102
Lyme disease, 102
rice-free,
Armigeres subalbatus, 108
ecology, 108
Japanese encephalitis virus, 108
Taiwan, 108
rickettsia,
Ixodes ricinus, 707
spotted fever group, 707
Rift Valley fever virus, 435, 467
mosquitoes, 467
Phlebotomus duboscqi, 435
Phlebotomus papatasi, 435
Phlebotomus sergenti, 435
reverse transcription-polymerase chain reaction, 467
Sergentomyia schwetzi, 435
vector competency, 435
Rio Grande do Norte,
leishmaniasis, 162
Lutzomyia spp., 162
sand flies, 162
Ross River virus, 146, 380
Aedes notoscriptus, 380
Australia, 146, 380
Barmah Forest virus, 380
dispersal, 380
mark-release-recapture, 380
mosquitoes, 146
vector competence, 146

S

salicylic acid,
6-methylsalicylic acid, 222
antifeedant, 222
Culicoides impunctatus, 222
repellent, 222
salicyluric acid, 222
salicyluric acid,
6-methylsalicylic acid, 222
antifeedant, 222
Culicoides impunctatus, 222
repellent, 222
salicylic acid, 222
salivary gland,
acquired resistance, 885
concanavalin A, 885
Ixodes scapularis, 885
lipopolysaccharide, 885
lymphocytes, 885
salivary gland immunity to ticks,
fecundity midgut, 601
feeding, 601
Rhipicephalus sanguineus, 601
tick performance, 601
salivary glands,
blood feeding, 743
hematophagy, 743
hyaluronidase, 743
Phlebotomus papatasi, 743
Simulium vittatum, 743
sample size,
altitude, 664
Anopheles, 664
epidemic, 664
forecasting methods, 664
malaria, 664
sampling, 299, 335
abundance, 299
accuracy, 299
bias, 299
Haematobia irritans, 299
*Ixodes ricinus*tick, 335
methodology, 335
precision, 299
sampling efficiency,
Ixodes scapularis, 357
Lyme disease, 357
population size, 357
tick density, 357
sand flies, 162, 289, 325
genetic variation, 325

isoenzymes, 289
isozymes, 325
leishmaniasis, 162
Lutzomyia longipalpis, 325
Lutzomyia spp., 162
Phlebotomus brevis, 289
Phlebotomus simici, 289
Rio Grande do Norte, 162
scanning electron microscopy, 289
Venezuela, 325
sand fly,
Lutzomyia, 860
sensilla coeloconica, 860
ultrastructure, 860
sand fly cells,
heterologous promoters, 802
luciferase expression, 802
Lutzomyia, 802
Phlebotomus, 802
sandflies,
Lutzomyia migonei, 748
mouthparts, 748
sarcophagid,
cutaneous myiasis, 854
scanning electron microscopy, 854
scanning electron microscopy, 289, 854
cutaneous myiasis, 854
isoenzymes, 289
Phlebotomus brevis, 289
Phlebotomus simici, 289
sand flies, 289
sarcophagid, 854
scFv,
Anopheles gambiae, 528
Monoclonal antibody, 528
phage display, 528
screening, 528
transmission blocking, 528
scorpions,
Androctonus, 787
genetic diversity, 787
ITS, 787
ribosomal DNA, 787
screening,
Anopheles gambiae, 528
Monoclonal antibody, 528
phage display, 528
scFv, 528
transmission blocking, 528
season,
Borrelia, 9
diel activity, 9
litter, 9
temperature gradient, 9
tick, 9
seasonal abundance, 153, 675
Anopheles farauti, 153
Australia, 153
bluetongue virus, 675
Culicoides sonorensis, 675
malaria, 153
sibling species, 153
survivorship, 675
vectorial capacity, 675
self-treatment, 588, 878
acaricide, 588, 878
Amblyomma americanum, 588, 878
amitraz, 588, 878
lone star tick, 588, 878
sensilla coeloconica,
Lutzomyia, 860
sand fly, 860
ultrastructure, 860
Sergentomyia,
antenna, 575
Indian sand flies, 575
larvae, 575
mouthparts, 575
Phlebotomus, 575
Sergentomyia schwetzi,
Phlebotomus duboscqi, 435
Phlebotomus papatasi, 435
Phlebotomus sergenti, 435
Rift Valley fever virus, 435
vector competency, 435
seroprevalence,
cattle, 27
enzyme-linked immunosorbent assay, 27
Hypoderma spp., 27
Spain, 27
warble flies, 27
sex,
body weight, 198
Dermacentor variabilis, 198
development, 198
Ixodes scapularis, 198
tick, 198
sheep nose bot,
morphology, 239

Oestrus ovis, 239
parasite, 239
puparium, 239
sibling species,
Anopheles farauti, 153
Australia, 153
malaria, 153
seasonal abundance, 153
Simulium damnosum,
heteroduplex analysis, 547
mitochondrial genes, 547
Onchocerca volvulus, 547
phylogeny, 547
Sudan, 547
Simulium damnosum complex,
chromosomes, 29
hybridization, 29
laboratory rearing, 29
multiple mating, 29
Simulium venustum,
Aedes communis, 919
AI3-37220, 919
deet, 919
piperidine, 919
repellents, 919
Simulium vittatum,
blood feeding, 743
hematophagy, 743
hyaluronidase, 743
Phlebotomus papatasi, 743
salivary glands, 743
small mammals,
climate, 820
interrelationship, 820
Ixodes didelphidis, 820
Ixodes loricatus, 820
soil chemistry,
aquatic ecology, 58
aquatic insects, 58
biting midges, 58
vector distribution, 58
Spain,
cattle, 27
enzyme-linked immunosorbent assay, 27
Hypoderma spp., 27
seroprevalence, 27
warble flies, 27
spatial temporal distribution,
Anopheles aquasalis, 231
Anopheles oswaldoi, 231
environmental factors, 231
larval habitats, 231
malaria, 231
species composition,
Costa Rica, 216
diversity, 216
Leishmaniasis, 216
Lutzomyia, 216
spectrofluorometry,
Aedes polynesiensis, 769
age-grading, 769
Culex quinquefasciatus, 769
fluorescence, 769
pteridines, 769
spermathecal ducts,
aedeagal filaments, 653
arms race, 653
coadaptation, 653
Phlebotomine sand flies, 653
Spinturnicidae,
bat, 445
Indiana, 445
Myotis sodalis, 445
parasitic mite, 445
Paraspinturnix globosa, 445
spontaneous flight activity,
Culex tarsalis, 187
longevity, 187
western equine encephalomyelitis virus, 187
spotted fever group,
Ixodes ricinus, 707
rickettsia, 707
St. Louis encephalitis virus, 250, 259, 507, 626
antibody, 250, 259
arboviral transmission, 626
avian serology, 507
California, 507
Culex tarsalis, 250
house finch, 259
infection, 259
mosquito blood feeding, 250
mosquito-borne encephalitis, 626
surveillance, 507
vector infection rates, 626
viremia, 250, 259
western equine encephalomyelitis virus, 250, 259, 507
wetlands, 507
stable fly, 308, 957

attraction, 957
feces, 957
host-location behavior, 308
human odors, 308
Stomoxys calcitrans, 308, 957
triple cage olfactometer, 308, 957
standards,
assay, 571
Ctenocephalides felis, 571
development, 571
larvae, 571
starch grains,
Phlebotomus papatasi, 319
photosynthesis, 319
plant feeding, 319
sugars, 319
stereoisomers,
1-[3-cyclohexen-1-ylcarbonyl] piperidine, 182
1-[3-cyclohexen-1-ylcarbonyl]-2-methylpiperidine, 182
Anopheles stephensi, 182
repellents, 182
Stomoxys calcitrans, 308, 957
attraction, 957
feces, 957
host-location behavior, 308
human odors, 308
stable fly, 308, 957
triple cage olfactometer, 308, 957
subterranean,
Aedes aegypti, 846
arbovirus, 846
Australia, 846
Mesocyclops, 846
mosquito, 846
Sudan,
heteroduplex analysis, 547
mitochondrial genes, 547
Onchocerca volvulus, 547
phylogeny, 547
Simulium damnosum, 547
sugars,
Phlebotomus papatasi, 319
photosynthesis, 319
plant feeding, 319
starch grains, 319
surveillance,
avian serology, 507
California, 507
St. Louis encephalitis virus, 507
western equine encephalomyelitis virus, 507
wetlands, 507
survival,
adult, 968
cryopreservation, 968
primary screwworm, 968
pupa, 968
survivorship, 385, 675, 701
Aedes albopictus, 701
Anopheles quadrimaculatus, 701
bluetongue virus, 675
cannibalism, 385
Coquillettidia perturbans, 701
Culicoides sonorensis, 675
eastern equine encephalomyelitis virus, 701
life tables, 385
mortality, 385
seasonal abundance, 675
Toxorhynchites rutilus, 385
vectorial capacity, 675
Sus,
host, 640
Manis, 640
parasite, 640
Thailand, 640
tick, 640
susceptibility, 194, 286, 331, 780
Anopheles albimanus, 331
Anopheles arabiensis, 286
Anopheles gambiae complex, 286
Culex pipiens molestus, 780
Culex tarsalis, 194
Culex tritaeniorhynchus, 780
DDT, 286
Iowa, 194
Japanese encephalitis virus, 780
multiplication, 194
phenotypes, 331
Plasmodium vivax, 331
synthetic pyrethroids, 286
Taiwan, 780
transmission, 194
western equine encephalomyelitis virus, 194
synergism,
cross-resistance, 721
insecticide resistance, 721
Pediculus capitis, 721
synthetic pyrethroids,

Anopheles arabiensis, 286
Anopheles gambiae complex, 286
DDT, 286
susceptibility, 286
systematic,
biting midges, 65
bluetongue, 65
Culicoides variipennis complex, 65
vectors, 65

T

Taenia solium,
flies, 489
Mexico, 489
Musca domestica, 489
vectors, 489
Taiwan, 108, 780
Armigeres subalbatus, 108
Culex pipiens molestus, 780
Culex tritaeniorhynchus, 780
ecology, 108
Japanese encephalitis virus, 108, 780
rice-free, 108
susceptibility, 780
Tanzania,
Culex quinquefasciatus, 172
gravid trap, 172
temperature,
distribution, 205
Forcipomyia taiwana, 205
population abundance, 205
rainfall, 205
temperature gradient,
Borrelia, 9
diel activity, 9
litter, 9
season, 9
tick, 9
teratology,
anomalies, 807
embryonic development, 807
Hyalomma marginatum, 807
Thailand, 77, 89, 640
Aedes aegypti, 77, 89
histology, 89
host, 640
Manis, 640
multiple blood feeding, 89
parasite, 640
population dynamics, 77
Puerto Rico, 77, 89
Sus, 640
tick, 640
tick, 9, 198, 640
body weight, 198
Borrelia, 9
Dermacentor variabilis, 198
development, 198
diel activity, 9
host, 640
Ixodes scapularis, 198
litter, 9
Manis, 640
parasite, 640
season, 9
sex, 198
Sus, 640
temperature gradient, 9
Thailand, 640
tick control,
ivermectin, 126
Ixodes scapularis, 126
Lyme disease, 126
white-tailed deer, 126
tick density,
Ixodes scapularis, 357
Lyme disease, 357
population size, 357
sampling efficiency, 357
tick ecology,
Amblyomma americanum, 439
Amblyomma maculatum, 439
Dermacentor variabilis, 439
Ixodes scapularis, 439
tick performance, 71, 601
fecundity immunity to ticks, 71
fecundity midgut, 601
feeding, 71, 601
Rhipicephalus sanguineus, 71, 601
salivary gland immunity to ticks, 601
ticks, 472, 774, 777, 973, 979
Amblyomma cajennense, 774
Amblyomma oblongoguttatum, 774, 777
Amblyomma ovale, 774
arrestant, 472
avoidance, 472
Brazil, 774
deer, 472
dogs, 774
ectoparasites, 973
Guinea-Bissau, 973
gynandromorphism, 777

hosts, 973
Ixodes scapularis, 472
Ixodidae, 973
kairomone, 472
Pantanal region, 979
wild animals, 979
Toxorhynchites rutilus,
cannibalism, 385
life tables, 385
mortality, 385
survivorship, 385
transmission, 114, 194, 835
adaptations, 114
behavior, 114
Culex tarsalis, 194
Dermacentor albipictus, 114
host-finding, 114
Iowa, 194
larvae, 114
mosquitoes, 835
multiplication, 194
Peru, 835
susceptibility, 194
vector competence, 835
Venezuelan equine encephalomyelitis virus, 835
western equine encephalomyelitis virus, 194
transmission blocking,
Anopheles gambiae, 528
Monoclonal antibody, 528
phage display, 528
scFv, 528
screening, 528
trematode vector,
aquatic insect host, 619
Ehrlichia risticii, 619
Potomac horse fever, 619
Triatoma,
fecundity, 373
feeding source, 373
life cycle, 373
Triatoma brasiliensis,
Chagas disease, 872
isoenzymes, 872
morphometrics, 872
population genetics, 872
random amplified polymorphic DNA, 872
Triatoma virus,
biological control, 271
insect picornaviruses, 271
triatominae,
Dipetalogaster maximus, 938
fatty acid transport, 938
fatty acid-binding protein, 938
muscles, 938
triple cage olfactometer, 308, 957
attraction, 957
feces, 957
host-location behavior, 308
human odors, 308
stable fly, 308, 957
Stomoxys calcitrans, 308, 957
tsetse,
Glossina morsitans centralis, 897
Glossina morsitans morsitans, 897
Glossina swynnertoni, 897
hybridization asymmetry, 897
turkey disease,
coronavirus, 480
darkling beetle, 480
lesser mealworm, 480
litter beetle, 480
poult enteritis mortality syndrome, 480

U

ultra-low volume insecticides,
Aedes aegypti, 541
culicidae, 541
resting behavior, 541
ultrastructure,
Lutzomyia, 860
sand fly, 860
sensilla coeloconica, 860

V

vector competence, 146, 695, 766, 784, 835
Aedes sollicitans, 695
Aedes taeniorhynchus, 695
Australia, 146
biting midge, 784
Borrelia burgdorferi, 766
Culicoides sonorensis, 784
Dirofilaria immiti, 695
dog heartworm, 695
Ixodes muris, 766
mosquitoes, 146, 695, 835
oral susceptibility, 784
Peru, 835

Ross River virus, 146
transmission, 835
Venezuelan equine encephalomyelitis virus, 835
vesicular stomatitis virus, 784
vector competency,
Phlebotomus duboscqi, 435
Phlebotomus papatasi, 435
Phlebotomus sergenti, 435
Rift Valley fever virus, 435
Sergentomyia schwetzi, 435
vector distribution,
aquatic ecology, 58
aquatic insects, 58
biting midges, 58
soil chemistry, 58
vector infection rates,
arboviral transmission, 626
mosquito-borne encephalitis, 626
St. Louis encephalitis virus, 626
vectorial capacity,
bluetongue virus, 675
Culicoides sonorensis, 675
seasonal abundance, 675
survivorship, 675
vectors, 65, 295, 489
biting midges, 65
bluetongue, 65
Culicoides variipennis complex, 65
Culiseta incidens, 295
Dirofilaria immitis, 295
flies, 489
heartworm, 295
Los Angeles County, 295
Mexico, 489
Musca domestica, 489
systematic, 65
Taenia solium, 489
Venezuela,
genetic variation, 325
isozymes, 325
Lutzomyia longipalpis, 325
sand flies, 325
Venezuelan equine encephalomyelitis virus,
mosquitoes, 835
Peru, 835
transmission, 835
vector competence, 835
vesicular stomatitis virus,
biting midge, 784
Culicoides sonorensis, 784
oral susceptibility, 784
vector competence, 784
viremia, 250, 259
antibody, 250, 259
Culex tarsalis, 250
house finch, 259
infection, 259
mosquito blood feeding, 250
St. Louis encephalitis virus, 250, 259
western equine encephalomyelitis virus, 250, 259
virulence,
arbovirus infection, 815
Culiseta melanura, 815
eastern equine encephalomyelitis virus, 815
mosquito mortality, 815
virus isolation methods,
Aedes aegypti, 641
antibody effect, 641
anti-dengue serum, 641
dengue virus, 641

W

warble flies,
cattle, 27
enzyme-linked immunosorbent assay, 27
Hypoderma spp., 27
seroprevalence, 27
Spain, 27
weather,
Ixodes scapularis, 408
Lyme disease, 408
population dynamics, 408
white-footed mice, 408
western equine encephalomyelitis virus, 187, 194, 250, 259, 507
antibody, 250, 259
avian serology, 507
California, 507
Culex tarsalis, 187, 194, 250
house finch, 259
infection, 259
Iowa, 194
longevity, 187
mosquito blood feeding, 250
multiplication, 194
spontaneous flight activity, 187
St. Louis encephalitis virus, 250, 259, 507

surveillance, 507
susceptibility, 194
transmission, 194
viremia, 250, 259
wetlands, 507
wetlands,
avian serology, 507
California, 507
St. Louis encephalitis virus, 507
surveillance, 507
western equine encephalomyelitis virus, 507
white-footed mice,
Ixodes scapularis, 408
Lyme disease, 408
population dynamics, 408
weather, 408
white-tailed deer,
ivermectin, 126
Ixodes scapularis, 126
Lyme disease, 126
tick control, 126
wild animals,
Pantanal region, 979
ticks, 979
Wolbachia, 53, 340
Armigeres subalbatus, 53
classification, 340
cytoplasmic incompatibility, 53
endosymbiont, 340
mosquito, 340
polymerase chain reaction, 340
wsp, 340
woodrats,
birds, 909
Borrelia burgdorferi, 909
Ixodes pacificus, 909
mice, 909
wsp,
classification, 340
endosymbiont, 340
mosquito, 340
polymerase chain reaction, 340
Wolbachia, 340

Y

Yersinia pestis,
fleas, 422
quantitative competitive polymerase chain reaction, 422
yolk deposition,
Anopheles stephensi, 726
Culex pipiens molestus, 726
Culex pipiens pallens, 726
fatty acids, 726
lipids, 726